

HYPERTENSION: TAKING THE PRESSURE OFF

By John Clark M.D. NorthernLightsHealthEducation.com

NATIONAL WATER SHORTAGE

“My eighty-three-year-old mother called me last week: ‘Brenda!’ her voice sounded alarmed, ‘I just took my blood pressure and I am worried; it is 160/100’”

Brenda was worried, too, but knew something about blood pressure. “Mom, just sit down, drink 3 big glasses of water, and I’ll be over in about 45 minutes and we’ll take your blood pressure again.” Arriving 45 minutes later, Brenda retook mom’s blood pressure; it was now 130/70.

Americans are chronically dehydrated. People who drink plenty of water have lower blood pressures.¹ Many people suffer from high blood pressure, the cause of which is inadequate consumption of water. Constant dehydration forces the body to tighten the blood vessels and speed up the heart to maintain adequate blood flow to the brain and over time hypertension results.² Not all hypertension is the result of not drinking enough water, but more of it arises from this cause than is realized.

Contrast the impact of this “national water shortage” with the National Institute of Health’s (NIH) guidelines for treatment of hypertension.³ For instance, say your water intake has been dismally inadequate and your brain is crying for better blood supply. The body responds with a faster heartbeat and tighter blood vessels to squeeze the blood more effectively up to the top of your head.⁴ The doctor takes your blood pressure and, of course, discovers you have blood pressure that is higher than is considered safe. In line with practice guidelines. The first line treatment is a diuretic – a “water pill” – You go home, the dehydration is worse, compounded, and on a return trip to the doctor you are again discovered to be not only hypertensive, but your heart rate is up. Guidelines now recommend a beta-blocker. What does a beta-blocker do? Slows your fast heart rate. With that you go home, your heart is now slow, but the brain is still crying out for blood so all the blood vessels in your body get even tighter to squeeze the remaining blood to the top of your head.

Back to the doctor you go for follow up; low-and-behold the pressure is still up. The next recommended pill to fix the “resistant hypertension” is a calcium channel blocker. What

does a calcium channel blocker do? It relaxes all the blood vessel in your body. You mean the ones that tightened up to get the blood to the top of my head? Yes. Now I must hand it to the NIH, the first line therapy is really supposed to be lifestyle modifications, but in my estimation the modifications listed and the extent to which they are applied or encouraged lacks enthusiasm. Okay, so, I guess you can see that I am not “real big” on treating diseases you get from poor lifestyle practices with any other thing than correct lifestyle practices.

SAVE THE BRAIN

People with normal blood pressure live longer.^{5,6,7} What is more, their thinking is clearer. Studying the brains of people with high blood pressure, also called hypertension, researchers have discovered an association between hypertension, brain white matter defects, and difficulty thinking and remembering.⁸ People with high blood pressure develop white matter lesions in their brains at 10 times the rate of the normal population.⁹ Controlling blood pressure with medications does not stop brain deterioration. Some blood pressure medications make the brain deteriorate even faster.¹⁰ In order to stop brain deterioration, lifestyle habits responsible for both hypertension and dementia need to be addressed.

“SAFE” MAGIC POTIONS?

Some people are quick to look to alternative medicine for a magic potion to fix their lifestyle related diseases, thinking they will find a safer cure in supplements, herbs, or vitamins. Some of these over-the-counter pills contain agents with pharmacological action just like medications, but does it make any sense to take magic potions to treat disease while continuing to practice the lifestyle habits that cause the illness in the first place? It’s not that people have not found benefit from some of these pills. Agents with some evidence of benefit include coenzyme Q10, fish oil, garlic, vitamin C, L-Arginine,¹¹ grape seed extract,¹² quercetin,^{13,14} rosemary,¹⁵ cranberry,¹⁶ fennel,¹⁷ mistletoe,^{18,19} saffron,²⁰ hawthorn berry²¹ and valerian.²² But none of these pills change the

reasons or causes as to why your blood pressure went up in the first place.

NO SUBSTITUTION

I am not against pills just because they are pills, but some of these pills can have serious drawbacks. For example, people already on blood pressure pills respond more poorly to lifestyle changes.²³ They are somewhat trapped. What's more, Parkinson's disease is a side effect of some blood pressure medications.²⁴ What about Alzheimer's? People with low to normal blood pressure are protected from brain diseases like Alzheimer's.²⁵ Artificial low blood pressure, with medications, does not always avert progression to dementia.²⁶ People whose good lifestyles give them a healthy blood pressure are much more likely to be mentally acute as they get older.²⁷ I recommend making a healthy lifestyle your defense against dementia, not medications.

While medications are generally prescribed to address some disease from which a patient suffers, hypertension is a side effect of many medications. Oral contraceptives rise blood pressure 8 points on average.^{28,29} Antidepressant use doubles the risk for hypertension.³⁰ Anti-inflammatory drugs cause high blood pressure through kidney dysfunction.^{31,32,33} Acetaminophen (aka Tylenol) increases blood pressure³⁴ especially in patients with coronary artery disease (most Americans).³⁵

WHAT IS HYPERTENSION?

What is Hypertension? The National Institute of Health is clear on the classification of blood pressure. Blood pressure is recorded as systolic pressure over diastolic pressure. Systolic is a measure of the highest pressure the heart develops during its beat and diastolic is the lowest pressure when the heart is refilling with blood for another beat. Normal blood pressure goes no higher than 120/80. To go over 120/80, but no higher than 139/89 is to have pre-hypertension. Blood pressure over 139/89 is classified in two stages. Stage 1 hypertension is from 140-159/90-99. Any pressure of 160/100 or more is considered stage 2 hypertension.³ The higher your blood pressure the higher your chances of experiencing other life-threatening complications of hypertension.

People whose good lifestyles give them a healthy blood pressure are much more likely to be mentally acute (sharp/bright) as they get older.

GLOBAL IMPACT

Hypertension affects approximately 50 million people in the United States and 1 billion worldwide.³ Fully 29% of US adults have hypertension, around 68% are taking antihypertensive medication, but only 64% of those taking medication have their blood pressure controlled.³⁶ Nine out of ten 55-year-olds, in the United States, with normal blood pressure will develop hypertension before they die.³

TYPES OF HYPERTENSION

There are two major types of high blood pressure, primary and secondary.³⁷ Primary blood pressure is generally considered to have an unknown cause, but as you continue to learn the facts about hypertension you will discover that most primary hypertension is lifestyle related—it is caused by our habits relating to eating, drinking and exercising. Secondary hypertension can be linked to other medical diseases such as thyroid disease, kidney disease, parathyroid disease, metabolic syndrome (diabetes), etc.

High blood pressure is called “the silent killer” because the majority of people with hypertension are unaware of its presence.

THE SILENT KILLER

High blood pressure is called “the silent killer” because the majority of people with hypertension are unaware of its presence. Nevertheless, some people with hypertension do report symptoms, these can include tiredness, sudden hot flashes, headaches, reduced energy, heart beat palpitations, sudden sweating, reduced physical performance, dizziness, shortness of breath, chest pain, sleepiness, blurred vision, tinnitus (ringing in the ears), and/or muscle tension.³⁸

BE ALERT FOR SIGNS OF CARDIOVASCULAR DANGER

Hypertension might not be so bad if it were not for all the other diseases it causes. Of these heart disease tops the list. Achieving normal blood pressure reduces the risk of congestive heart failure by 36%,³⁹ of coronary heart disease by 72%, and of heart attack by 75%!⁴⁰ Because hypertension is so hard on the heart, if you can reduce the blood pressure by just 20 points, you can cut the risk of heart disease in half.³

Researchers have determined that lowering the average blood pressure in the United States by just 5 points would reduce the number of stroke deaths by 23,000 per year.⁴¹ Further, lowering your blood pressure from 140/90 to a more normal 120/75 reduces your risk of stroke by 74%.⁴² Bleeds are a common cause of stroke and can occur when an aneurysm bursts in the brain. If blood pressure is kept within normal limits, brain aneurysms are much less likely to enlarge and rupture.⁴³

Another study reported that normalizing blood pressure can reduce strokes by 35–40%, myocardial infarctions by 20–25% and heart failure by more than 50%.⁴⁴

Controlling blood pressure also reduces the risk of peripheral vascular disease – injury to blood vessels in your arms and legs that can lead to physical disability.⁴⁵

ADDING INSULT TO INJURY

Hypertension can damage any part of your body because every part of your body is dependent upon blood for life. People with normal blood pressure save themselves much grief.

People with normal blood pressure can avoid the second leading cause (behind diabetes) of kidney failure.⁴⁶

A surprising consequence of high blood pressure is high cancer risk! For example, normal blood pressure decreases the risk of endometrial cancer by 70%.⁴⁷

Do you remember your last blood pressure numbers? Are you having difficulty thinking and remembering? People with hypertension are more apt to get Alzheimer's in their later years, a disease of the brain that affects the ability to think and remember. People with normal blood pressure have superior blood flow to their brains improving thought and memory.^{48,49} Normal blood pressure protects you from brain deterioration especially in the frontal lobes.⁵⁰ People who do not have hypertension perform better on test of memory, attention, and abstract reasoning.⁵¹ Normal blood pressure can actually reduce the risk of Alzheimer's by 40%.⁵²

Diabetes and hypertension are often closely related. When combined with low HDL cholesterol, high triglycerides, and central obesity they are given a diagnosis of syndrome X.⁵³ Syndrome X is not a good diagnosis to have, but lifestyle interventions are most effective at treating this disease.

Osteoporosis is a thinning of the bones leading to an increased risk of fractures. People with normal blood pressure have 1/3 the risk of osteoporosis as those with hypertension.⁵⁴

Retinopathy and Macular Degeneration are leading causes of blindness. Keeping blood pressure within normal limits cuts the risk of blindness from retinopathy and/or macular degeneration in half.⁵⁵

Erectile dysfunction; the pressure in performance anxiety! Having trouble pleasing your wife? Thirty-two percent more men with normal blood pressure “get it up” than men with hypertension.⁵⁶

WHERE ARE WE HEADED: OUTLINE

Where we are headed with our hypertension discussion? Blood pressure is a product of the pumping of the heart, the blood vessel size, blood thickness and blood volume. If the heart beats faster, more blood is pumped and the pressure goes up. We call this tachycardia. If the blood vessels tighten up, making the space for the passage of blood narrower, it takes greater pressure to get the same amount of blood through to its destination. We refer to this tightening up effect as vasoconstriction. Blood vessels normally expand with each beat of the heart and then relax. If the blood vessels become hard, their stiffness inhibits the free flow of blood with each beat of the heart and the pressure goes up. Atherosclerosis is an example of this process. If the blood becomes thick and sludgy, more pressure is required to push it through the blood vessels and hypertension commences. When blood gets thick, we say that the viscosity has increased too much. If the blood vessels are being choked by something pressing on them from their sides, the effect is like putting your thumb over the end of a garden hose; this results in increased blood pressure. We call this external compression. Finally, if the volume of blood increases, this increases the amount of blood entering the heart, this in turn increases the amount of blood the heart pumps with each beat increasing the overall blood pressure. We often refer to this phenomenon as fluid retention.

VOLUME OVERLOAD!

Let us start by talking about the problem of volume overload. Volume overload can be the result of obesity, where it takes more blood to feed an increased mass of fat tissue. Salt causes fluid retention effectively producing volume overload. Volume overload results when the kidneys fail, because the kidneys are responsible for dispensing with excess fluid volume. Muscles have many large blood vessels and require lots of blood when exercised. Big muscles left unused become stiff and their blood vessels become stiff resisting blood flow and causing volume overload at the heart. Thus,

inactivity leads to hypertension. Blood does not like to be cold (you are not cold-blooded) and if your legs or arms are cold, the blood vessels in these extremities tighten up, sending all of the blood flooding to the heart, overloading it and causing hypertension.⁵⁷

US SODIUM INTAKE EXCEEDS DIETARY GUIDELINES

It is a well-known fact that dietary salt plays a significant role in the evolution of hypertension. Salt causes your body to hold on to excess fluid, causing volume overload at the heart, and subsequent hypertension. It is estimated that reducing sodium intake in America to 1300mg (about ½ teaspoon of salt) per day would reduce the yearly death rate by 150,000.⁵⁸ Our recommends are that total salt consumption not exceed 1/4 to 1/8 teaspoon per day, or 600mg to 300mg of sodium total.

Salt is a popular ingredient in fast food restaurant menu items. Salt and hypertension are key ingredients for the occurrence of a brain stroke. A national statistic reveals that the more neighborhood fast food restaurants a community has, the higher will be the stroke rate in that given community.⁵⁹

Okay, so test your salt savvy; which has more sodium – Rice Chex cereal or potato chips per one ounce serving each? The Rice Chex cereal, at 249 mg, has nearly twice as much as potato chips at 147 mg. How did you do? Now, compare tortilla chips and canned tomato sauce. The tomato sauce tops the chips at 147 mg / oz, the chips contain 118mg of sodium per ounce. What about Kraft Velveeta processed cheese or a hot dog sandwich? Ounce for ounce the Velveeta has nearly twice as much sodium at 420 mg compared to the hot dog's 221 mg.⁶⁰ Read your labels! Don't get caught off guard. Or, better yet, buy food without labels, like corn on the cob, which has one-tenth the sodium as canned corn.⁶¹ In Japan, soy sauce is a significant source of excess sodium. In Japan, most (63%) dietary sodium comes from soy sauce and is a big cause of hypertension.⁶²

Which has more sodium Rice Chex cereal or potato chips per one ounce serving each? The Rice Chex cereal at 249 mg has nearly twice as much as potato chips at 147 mg.

Let us suppose your sodium intake is within safe limits, but you still seem to be having salt related hypertension. It may be due to other things in your diet that cause you to retain sodium. Refined

carbohydrates and saturated fats increase salt retention and lead to hypertension.^{63,64}

Psycho-social factors can also affect the body's propensity to hold on to excess sodium. People under stress retain sodium.⁶⁵ Thus, people who don't stress out, retain less salt and have lower blood pressures. We will be revisiting this factor in detail under the fast heart rate discussion section further on in this book. Blood-pressure-raising-sodium is not unique (or limited) to table salt; it also appears in sea salt and in mono sodium glutamate (MSG). It should come as no surprise then that MSG consumption increases the likelihood that a person will get high blood pressure.⁶⁶

ESCALATING OBESITY RAISES PRESSURE CONCERNS

Think twice before shopping for a larger dress; clothing size correlates directly with increases in blood pressure.⁶⁷ According to the World Health Organization, more than one billion people worldwide are overweight and more than 300 million people are obese resulting in high rates of hypertension, kidney disease and cardiovascular disease.⁶⁸ What we eat in America, the (S)tandard (A)merican (D)iet, makes the liver sick with "fatty liver disease" and triples the risk of hypertension.⁶⁹ A "pot belly" is a bad omen for hypertension. Thinner is better, tighten that belly for a drop in pressure! The more abdominal (visceral fat) you sport, the greater your risk of hypertension.⁷⁰ One serving of cheese per day can significantly increase a man's waist circumference, body mass index and blood pressure.⁷¹ Indeed, each inch you can tighten your belt lowers your risk of hypertension by 15%.⁷² To relate it to absolute weight gain in pounds, a 55 lb weight gain over your ideal body weight raises your risk of hypertension by 265%.⁷³ It is estimated that in up to 50% of the adults in the United States, whose hypertension is being managed with pills, the need for drug therapy could be alleviated with modest reductions in body weight.⁷⁴

INEQUALITY AMONG THE PROTEIN GIANTS

While many see protein as essential, excessive intake has been linked to hypertension.⁷⁵ Specifically, animal protein especially decreases kidney function increasing the risk of hypertension.⁷⁶ On the other hand plant protein has been demonstrated to lower blood pressure. Increased intake of plant protein, fruits and vegetables significantly lowers the risk of hypertension.⁷⁷ As a practical example, two groups of people were compared. The first group used milk,

a source of animal protein, and the second group was given soy milk, a source of vegetable protein. Those on the soy milk experienced 18 mmHg lower blood pressures than those on the cow's milk.⁷⁸ What's more a switch to soy can improve kidney function and insulin sensitivity,⁷⁹ and lower serum total cholesterol levels.⁸⁰

VASOCONSTRICTION

Let's now turn our attention to the impact of vasoconstriction on the development of high blood pressure. If the blood vessels tighten up, making the space for the passage of blood narrower, it takes greater pressure to get the same amount of blood through to its destination. What makes the blood vessels tighten up? Psychological stress, cold – especially in the arms and/or legs, and failing to maintain adequate water intake. Then there are the substances which stimulate the blood vessels to constrict such as caffeine from tea, coffee, colas, etc, and tobacco.

PUTTING PRESSURE ON: SUBSTANCE ABUSE

Caffeine makes the heart react as it would if you were in a real life-or-death, stressful, flight-or-fight situation.⁸¹ What actually happens is that caffeine acutely raises blood pressure by raising circulating concentrations of the stress mediators epinephrine and norepinephrine. In addition, caffeine increases arterial stiffness and inhibits the relaxation of blood vessels.⁸² The impact of caffeine consumption on blood pressure is dose dependant – the more caffeine you consume, the more your blood pressure increases.⁸³

Chocolate contains caffeine as well as other similar vasoactive substances such as theobromine.⁸⁴ Some have crafted studies (which lack any disclaimer to industry involvement, funding, or researcher bias) designed to give chocolate apparent positive effects on high blood pressure.⁸⁵ But in studies of real people eating chocolate available from stores it does not lower blood pressure, it only tends to encourage eating between meals and weight gain.⁸⁶

The nicotine in tobacco is also a vasoconstrictor and pressor—a substance which raises blood pressure. Non-smokers have 12% lower risk of developing hypertension than smokers.⁸⁷

ENVIRONMENTAL HAZARDS

Remember the people who got sick from the FEMA trailers used to house victims of hurricane Katrina that hit Louisiana?⁸⁸ Environmental

chemicals such as formaldehyde and acetaldehyde, which are found in building materials and cigarette smoke, increase hypertension, tightening blood vessels and increasing the amount of blood the heart pumps.⁸⁹

Vasoconstriction can be the result of electromagnetic bombardment. For example, 40 minutes on the mobile phone can raise your blood pressure by 10 points.⁹⁰

WEATHER AND CLOTHING

When your arms or legs get cold, the blood vessels in them tighten up to reduce the amount of blood coming to them so that you will not lose too much heat. Poorly clad, chilled extremities force blood back to the heart, doubling its work and raising blood pressure.⁹¹ It is interesting to note that blood pressure increases in the winter, especially in the elderly,⁹² but so does consumption of salt and fat.⁹³

On the brighter side, don't underestimate the benefits of sunshine! Sunshine relaxes blood vessels lowering blood pressure^{94,95} and increases vitamin D, which has also been shown to lower blood pressure.⁹⁶

MOVE THOSE MUSCLES!

Don't take life sitting down! Active people have lower blood pressures; sedentary ones get hypertension.⁹⁷ Inactivity leads to increased vascular resistance to blood flow,⁹⁸ decreased blood flow to large muscles,⁹⁹ and increased blood pressure.¹⁰⁰ Regular use of your muscles keeps them supple and well supplied with blood, this in turn lowers blood pressure.^{101,102}

The benefits of exercise in the treatment of hypertension are often overlooked. Exercise is important for all aspects of health. If you keep wiggling, they won't put you in a box! As a mode of exercise, walking is hard to beat. In fact, walking 10,000 steps or more per day can lower your blood pressure by 10 points.¹⁰³ That would be about 4 miles. Weight lifting or resistance training can provide additional benefit. Indeed, 20 minutes per day in the gym can lower you blood pressure by as much as 10-12 points.¹⁰⁴

RELAXATION

Feel the need of a massage? A back massage brings relaxation and lowers blood pressure.¹⁰⁵ A nice soothing warm bath is also beneficial.¹⁰⁶

EXTERNAL COMPRESSION

External compression, the choking off of blood flow by something pushing on the blood vessel from its outside, causes hypertension. The effect is like putting your thumb over the end of a garden hose; the result is increased blood pressure. If someone grabs your neck from behind and chokes off your air and carotid arteries, not only will your eyes bulge, but your blood pressure will probably go up. Physical things which produce an external compression of the blood vessels include swelling, or edema, inflammation, sugar coating of the vessel walls called glycation, tight clothing, and obesity.

TIGHT CLOTHING

Tight clothing, like belts and elastic, compress blood vessels and raise blood pressure. Clothing that hangs from your shoulders leaves your waist free of compression and aids in lowering blood pressure. Wearing loose clothes allow for more free blood flow and more normal blood pressure.¹⁰⁷ Instead of wearing a belt to hold up pants, suspenders are helpful to avoid the tightness.

BREATHE CORRECTLY

Deep abdominal breathing, in contrast to shallow chest/neck breathing, brings blood pressure down.^{108,109}

BLOOD VISCOSITY, THICK BLOOD

If the blood becomes thick and sludgy, more pressure is required to carry it through the blood vessels and hypertension commences.¹¹⁰ When blood thickens, we say that the viscosity has increased too much. What happens is that the red blood cells stick together in a series or chain. We call this phenomenon rouleaux.¹¹¹ Factors known to influence blood toward increased viscosity and rouleaux include stress, dehydration, consuming refined foods, high blood cholesterol, high fat diet, overeating, and the accumulation of waste products in the intestines.

Refined foods, such as oils and sugars, make red blood cells stick together in clusters or chains called rouleaux that can be seen under a microscope. Higher blood pressure is required to circulate this thickened blood.¹¹²

FATS

Because eating a high amount of fat causes rouleaux, food with saturated fat, (i.e. animal

products, butter,¹¹³ margarine, shortening, and/or coconut oil,¹¹⁴ etc.) significantly increases the risk of hypertension. In addition, consumption of these products also raises the blood cholesterol, thickening the blood further, and raising blood pressure.¹¹⁵ Even God has weighed in on the fat question, "Speak unto the children of Israel, saying, Ye shall eat no manner of fat..."¹¹⁶

Do you know how much fat is in the food you eat? Fat in the diet increases blood pressure because it thickens the blood, and makes the blood vessels stiff.

Compared to naturally occurring vegetable fats, lard significantly increases blood pressure and risk of hypertension.¹¹⁷ This problem is worse as one gets older.¹¹⁸ These scientific facts help explain why God said, "And the swine, because it divideth the hoof, yet cheweth not the cud, it is unclean unto you: ye shall not eat of their flesh, nor touch their dead carcase."¹¹⁹ One group of researchers wanted to see what would happen if people were placed on an animal fat free diet for 10 days. It was discovered that ten days on an animal fat free diet significantly reduced blood pressure.¹²⁰ It is interesting to note that a similar experiment was entered upon around 600 B.C. in Babylon.¹²¹ Daniel was a Jewish captive of Babylon. When offered a diet of meat he requested, "Prove thy servants, I beseech thee, ten days; and let them give us pulse (vegetables) to eat, and water to drink."¹²² The outcome was that Daniel and three other colleagues who shared the intervention arm of the study did ten times better in school than all the other university students.

What about processed vegetable oils? These oils are not free from health risk just because they come from plants. Refined oils, especially canola oil,¹²³ increase hypertension¹²⁴ and the risk of stroke at a younger age.¹²⁵

Another problem with oils is what happens to them when they are subjected to heat. For example, heating oils in a pan causes serious deterioration in their quality turning them toxic so that they contribute to the onset and severity of hypertension.¹²⁶

Do you know how much fat is in the food you eat? Fat in the diet increases blood pressure because it thickens the blood, makes the blood vessels stiff,¹²⁷ and causes endothelial dysfunction.¹²⁸ What is endothelial dysfunction? Nitric oxide is used in the body to relax blood vessels. When the blood vessels respond poorly to nitric oxide relaxation we say it is a result of the inner lining or endothelium of the blood vessel being sick or dysfunctional, hence, endothelial dysfunction.¹²⁹ Foods that decrease

vascular sensitivity to nitric oxide relaxation signals include: high fat,¹³⁰ salt,^{131,132} cholesterol,^{133,134} (especially dietary oxidized cholesterol¹³⁵ ¹³⁶), overeating,^{137,138} sugar,^{139,140} (especially fructose^{141,142}), and glycation of proteins as happens in diabetes.¹⁴³

THE VASO-RELAXING DIET

On the other hand, proper diet has a significant impact on the responsiveness of your blood vessels to nitric oxide relaxation. I call it the vaso-relaxing diet, meaning it MSKRD your blood vessels more responsive to relaxation messages from your body. Dietary changes known to improve vascular responsiveness include: a vegetarian diet,¹⁴⁴ oats (oatmeal),¹⁴⁵ tomatoes,¹⁴⁶ diets rich in antioxidants¹⁴⁷ such as vitamin E,¹⁴⁸ and minerals like zinc¹⁴⁹ and copper.¹⁵⁰

Omega-3 deficiency leads to hypertension.¹⁵¹ Flaxseed and walnuts are good dietary sources of omega-3 fatty acids. Omega-3 fatty acids are very antithrombotic and anti-inflammatory. In contrast, omega-6 fatty acids, which are present in refined vegetable oils and meat, are prothrombotic (causing blood clots) and proinflammatory. Omega-3 fatty acids also aid in the treatment of hyperlipidemia, hypertension, and rheumatoid arthritis.¹⁵²

THE DEADLY MIX: FAT AND SUGAR

When combined, fat and sugar form a deadly mix. The risk of hypertension from combining these two agents is not just additive, it is multiplicative,¹⁵³ meaning just small amounts of these two agents mixed together creates an enormous health hazard.

THE AMERICAN SWEET TOOTH

Speaking of sugar, (and refined carbohydrates, like white flour, white pasta, white rice, and refined breakfast cereals, which turn immediately to sugar in your blood stream), 20 teaspoons of sugar raises your blood pressure by 2 mmHg, 40 teaspoons raises it by 5 mmHg.¹⁵⁴ The average American consumes 47 tsp of sugar each day of their lives.¹⁵⁵ The danger of sugar in the diet is that it ends up in the blood stream. As the blood sugar rises so does the blood pressure.¹⁵⁶

Not all carbohydrates are created equal. Sugar and complex carbohydrates may carry the same theoretical calories per gram, but the impact on the blood sugar, the risk of hypertension and diabetes complications are very different.¹⁵⁷ Complex carbohydrates from an unrefined plant-based diet are much better tolerated, decrease the risk of

hypertension, and provide a nutrient dense diet that improves health.¹⁵⁸ So how do sugar and other refined carbohydrates cause hypertension? Refined carbohydrates cause hypertension by increased production of norepinephrine, dopamine, and epinephrine, and by causing blood vessel wall thickening.¹⁵⁹ The secret is to quit consuming sugar; it works, and blood pressure comes down.¹⁶⁰

Sugar is an accumulative poison. Refined carbohydrates create a sticky coating of sugar all over your blood cells, blood vessels and other body tissues. The more refined processed foods you eat, the higher your blood sugar goes and the more this sticky sweet substance coats the tissues of your body. This coating is called glycation and accumulates over time, leading to vascular and heart muscle stiffness, atherosclerotic plaque and hypertension. The more refined food you eat in your lifetime the greater your chance of having high blood pressure as you age.¹⁶¹ People on an unrefined plant-based diet enjoy a cleaner cardiovascular system and may avoid high blood pressure all together as they get older.¹⁶²

Eating refined carbohydrates has a downside of leading to a condition called insulin resistance, where the cells of the body no longer take sugar out of the blood stream in response to normal insulin levels.^{163,164} Examples of refined carbohydrates shown to increase insulin resistance include: white rice,¹⁶⁵ white flour, as found in pastries and white bread,^{166,167} and processed sugar.^{168,169} Of particular concern among refined carbohydrates is fructose, its ability to create insulin resistance and hypertension surpasses table sugar.^{170,171}

Sugar is an accumulative poison. Refined carbohydrates create a sticky coating of sugar all over your blood cells, blood vessels and other body tissues. The more refined processed foods you eat, the higher your blood sugar goes.

Additional causes of insulin resistance include a high fat diet,¹⁷² eating between meals or snacking¹⁷³ and obesity.¹⁷⁴ The results of insulin resistance are high cholesterol values¹⁷⁵ and, of course, hypertension.^{176,177,178} People who never provoke their insulin to overproduction by eating refined carbohydrates also never experience insulin-driven hypertension.¹⁷⁹ Eat only as much refined carbohydrates as you would like to see your blood pressure go high.

Speaking of fructose, drinking one fructose-sweetened soda per day can increase the risk of hypertension by 77%.¹⁸⁰ And, in case you were

thinking artificial sweeteners were a good alternative, think again. Artificial sweeteners like aspartame give some people very high blood pressure.¹⁸¹ Stopping these sweeteners can be the secret to lowering some people's blood pressure.¹⁸² On the other hand there is a non-sugar sweetener which has been shown to improve blood pressure, and that is stevia.¹⁸³ It does this by acting as a calcium channel blocker (the mechanism of action of some pharmaceutical blood pressure drugs).¹⁸⁴

HARDENING OF THE ARTERIES

Hardening of the arteries causes resistance to free flow of the blood. Blood vessels normally expand and relax with each beat of the heart. If the blood vessels become hard, their stiffness inhibits the free flow of blood with each beat of the heart and the pressure goes up.^{185,186} Atherosclerotic plaque is an example of this process.¹⁸⁷ If a plaque narrows the diameter of a blood vessel, more pressure is required to get the same amount of blood through the smaller opening.

Other conditions leading to this mechanism of blood pressure elevation include; increased body inflammation¹⁸⁸ (also known as oxidative stress), physical inactivity, and endothelial dysfunction. Endothelial dysfunction is a deadening of the blood vessel wall so that it no longer responds to the bodies relaxing signals. Endothelial dysfunction leads to unresponsive arteries and veins. Unresponsive arteries and veins resist free flow of blood to the tissues raising the blood pressure required to move the same volume of blood. Things that cause endothelial dysfunction include high blood sugars with sugar coating of the endothelium, uric acid with deadening of the endothelium, and high fat which also deadens the endothelium so that it will not relax and allow blood to pass freely.

INFLAMMATION AND ENDOTHELIAL DYSFUNCTION

The role of inflammation in the development of hypertension is multifaceted. Besides causing endothelial dysfunction, it thickens blood vessel walls making blood flow more difficult and raising the pressure required for normal circulation. In consequence, blood pressure goes up with increasing inflammation.¹⁸⁹ People with more inflammation have a 40% higher risk of hypertension.¹⁹⁰ Many people realize that there is an association between salt intake and a rise in blood pressure. In fact, even though arterial hypertension is a major cause of disease-related morbidity and mortality worldwide, it is nearly

absent in populations that consume natural foods containing little or no salt.¹⁹¹ This is partly due to fluid retention,^{192,193} but it is also due to inflammation.¹⁹⁴ Excessive salt intake causes hypertension and kidney injury, in part, by oxidative stress that inflames the blood vessel walls and kidneys.¹⁹⁵

TOXINS IN THE ENVIRONMENT

Lead¹⁹⁶ and arsenic¹⁹⁷ are environmental poisons that raise whole body inflammation and have been shown to cause hypertension. Lead is a common contaminant in old paint, dust around old houses and can even be found in supplements such as calcium.¹⁹⁸ Arsenic often contaminates public drinking water¹⁹⁹ and bottled water.²⁰⁰ Arsenic is used in chicken and hog feed as a growth promoter and is present in the meats of these animals. Arsenic is an approved feed supplement that farmers use to control intestinal parasites in chickens.²⁰¹ As a consequence, eggs²⁰² and chicken meat²⁰³ prove to be dietary sources of arsenic. Seafood, because fish swim in polluted waters, has been found to be a significant source of arsenic in those that consume seafood.²⁰⁴ Avoid arsenic and avoid hypertension.

HOW MUCH ANTIOXIDANTS DO I NEED?

The secret to inflammation management is having a good offense. Antioxidants and phytochemicals from plant foods provide this; they lower the inflammation in your body^{205 206} and normalize your blood pressure.²⁰⁷ You may be asking; how much antioxidant do I need? Can I get too much? You may be interested to find out that lifestyle measures are not easily overdosed, they have more marked results in more difficult cases, and they do not cause dangerous side effects. Antioxidants only lower blood pressure in people with hypertension, not in normal healthy people.²⁰⁸

Diet is not the only way to have a positive impact on your antioxidant defense system. Exercise helps reduce the body's inflammation resulting in lower blood pressure.²⁰⁹

Since periodontal disease increases inflammation and hypertension, good oral hygiene can also be a key to good blood pressure control.²¹⁰

BEAUTY REST PRESCRIPTION

One of your bodies' own defenses against elevated inflammation is the hormone produced by the pineal gland in your brain, called melatonin. Melatonin is a natural antioxidant that reduces blood pressure.²¹¹ Late evening artificial lighting

impacts sleep, disrupts melatonin production, and raises blood pressure.^{212,213,214} People who take regular hours for adequate sleep in total darkness have a healthier supply of melatonin and lower blood pressures.^{215,216} Consequently, people who subject themselves to the irregularities in schedule afforded by shift work are at higher risk of hypertension.^{217,218} That extra night's pay may not be worth the risk of high blood pressure. Regular day jobs are friendlier to blood pressure. We recommend a 9:30 p.m. bedtime for adequate melatonin production and healthier blood pressure.

People who go to sleep easily and get plenty of rest experience less hypertension. People who breathe freely at night, without sleep apnea or snoring, have lower blood pressures.²¹⁹

MEAT AND ENDOTHELIAL DYSFUNCTION

So what raises inflammation increasing blood vessel wall thickness and causes endothelial dysfunction? Animal product consumption. Three or more servings of meat, poultry, eggs, fish, and/or seafood per day increases the risk of hypertension by 67%.²²⁰ Of particular concern are pork products. The American Heart Journal notes: "A complete elimination of pork in all forms is a good dietary therapeutic rule in the management of arterial hypertension in all patients." "The elimination of all forms of pork from the diet of all people may be a good practice for the prevention and the control of hypertension, one of the most common and important illnesses of man." "Salted pork is even more hazardous to the health of man."²²¹ Maybe this is why it was not considered a food item in God's book, for Leviticus notes, "And the swine...is unclean to you."²²² Even eggs won't help you lower your blood pressure. Eggs play a significant role in people 40 years old and older in their risk of hypertension.²²³ Animal products lack solution-oriented anti-oxidants, and if they are not a part of the solution, then they are often a part of the problem. This is definitely the case in eating animal products.

ANIMAL PRODUCTS FACE ACID TEST

Acid is no friend of your body's sensitive tissues. Acid increases endothelial dysfunction leading to hypertension.²²⁴ As a consequence, acid forming foods, particularly cheese and animal protein, significantly increase hypertension.²²⁵ Studying acid in the urine to identify acid forming foods, fruit and vegetables actually lower the amount of acid that is formed in the digestion of food. Grains, fish and red meat increase acid formation, but cheese almost

triples the amount of acid produced by comparison.²²⁶ One of the offending acids produced is uric acid. Anything that raises uric acid in the blood raises the risk of hypertension.²²⁷ Foods that raise uric acid include: meat, particularly organ meats,^{228,229} seafood,²³⁰ fructose,^{231,232} often as high fructose corn syrup or agave, beer and alcoholic beverages.²³³ As a consequence, the same diet helpful in lowering the risk of gout from uric acid elevations can simultaneously lower the risk of hypertension.

Foods known to help lower uric acid levels include fruits such as lemons²³⁴ and other citrus, cherries,²³⁵ strawberries,²³⁶ grapes, apples, tomatoes, bananas, and pomegranates; vegetables such as celery, potatoes, beets, and endive; and most tree nuts, especially chestnuts.²³⁷ For that matter, any food with diuretic properties has the potential to relieve high blood pressure that is being caused by high uric acid levels. Foods valued for their enhanced urine production properties (diuretic) include artichoke, celery, eggplant, cauliflower, green beans, grape, apple, peach, pear, melon, and watermelon.²³⁸ When you increase the amount of urine you produce, you potentially increase the amount of uric acid lost in the urine, thus lowering the levels in your blood.

STRONG DRINK

The impact of alcohol on the rise in blood pressure is not limited to its propensity to elevate blood uric acid levels. There is a linear relationship between the amount of alcohol consumed and the rise in blood pressure²³⁸ (the more alcohol you drink the higher your blood pressure goes²³⁹). A two drink a day person can lower their risk of hypertension by one third simply by abstaining.²³⁹ The wise man says; "Wine is a mocker, strong drink is raging: and whosoever is deceived thereby is not wise."^{240,241}

CHOOSING A BLOOD PRESSURE SENSITIVE DIETARY LIFESTYLE

Diet plays a significant role in hypertension, after all, you are what you eat! In a study comparing common diets, it was discovered that non-vegetarians, (consumers of animal products like meat, eggs, and dairy) eat 50% more fat, have 30% higher total cholesterol, have 32% higher blood sugars and are six times more likely to develop hypertension.²⁴² Indeed, in rural populations still practicing their traditional largely vegetable based lifestyle, free from the effects of the western diet, hypertension is unheard of and there is no age related rise in blood pressure.²⁴³ Their average

blood pressure is around 110/60 mmHg, average cholesterol 129 mg/dl, blood sugars 55 mg/dl and body mass index is 20 (kg/cm²).²⁴⁴

In a study comparing common diets, it was discovered that non-vegetarians (consumers of animal products like meat, eggs and dairy) are six times more likely to develop hypertension.

Really, blood pressure is not supposed to go up with age! In choosing a dietary lifestyle you will be interested to know that people adopting total plant based nutrition only have a 5% incidence of hypertension, those vegetarians including dairy and eggs with their meals, a 10% incidence, fish eating vegetarians a 12% incidence, and meat eaters a 21% incidence.²⁴⁵ Studied from another angle, people on total plant based nutrition have one fourth the risk of developing hypertension as meat eaters.²⁴⁶

In considering adopting a health promoting lifestyle, the vegan (plant-based nutrition) diet has many blood-pressure lowering advantages. In one study hypertensive patients, on blood pressure medications for 8 years, were given a vegan diet for one year. Blood pressures came down and the majority were able to stop or drastically reduce their medications.²⁴⁷ Just eating more fruits and vegetables is helpful, compared to those who do not eat many fruits or vegetables; those who eat largely of fruits and vegetables have a 77% lower risk of hypertension.²⁴⁸ Vegetables lower blood pressure and keep it from creeping up over the years.²⁴⁹

Vegetables known to be helpful in lowering blood pressure include: green leafy because they are high in magnesium and potassium,²⁵⁰ spinach because it is rich in folate,²⁵¹ celery^{252,253} (eat it or drink it), carrots,²⁵⁴ ginger,²⁵⁵ and broccoli,²⁵⁶ because they relax blood vessels, beets,²⁵⁷ basil,²⁵⁸ garlic^{259,260} (supplement or whole), onions; (eaten raw, but not cooked) significantly reduce hypertension.²⁶¹

Not to be left out, fruit also lowers blood pressure.²⁶² Fruits especially shown to have a positive impact on blood pressure include eggplant,^{263,264} tomatoes,²⁶⁵ pumpkin or squash²⁶⁶ (which are high in potassium and low in sodium), cranberry (which has ACE inhibitory activity mimicking a popular blood pressure pill),²⁶⁷ an apple a day,²⁶⁸ pomegranates,²⁶⁹ grapefruit²⁷⁰ (which protect arteries, has diuretic activity and improves blood fluidity), and finally olives²⁷¹ (which have calcium channel blocking and nitric oxide mediated vasodilatation benefits).

Deficiency of certain minerals in the diet increases the risk of hypertension. Iron: people

eating a nutritious vegetarian diet providing sufficient intake of iron enjoy lower blood pressures. Iron from meat, known as heme-iron, is not helpful in this regard.²⁷² Potassium: lowers blood pressure,²⁷³ but the most common blood pressure pill, a diuretic, lowers potassium.²⁷⁴ Calcium is needed to maintain normal blood pressure.²⁷⁵ Copper deficiency, a common result of eating refined carbohydrates,²⁷⁶ has been shown to cause high cholesterol (hypercholesterolemia), high triglycerides (hypertriglyceridemia), hypertension, and glucose intolerance (diabetes).²⁷⁷ Magnesium deficiency precipitates hypertension.²⁷⁸ Mineral levels are decreased by alcohol, salt, phosphoric acid (sodas), coffee intake, by profuse sweating, by intense prolonged stress, by excessive menstruation and vaginal flux, by diuretics and other drugs by certain parasites (pinworms), by sugar (refined carbohydrates), and by use of tobacco.

On a fresh fruit and vegetable diet, participants lost 8 pounds, lowered their blood pressures by 18 mm Hg, and 80% of those who smoked or drank alcohol abstained spontaneously!

WHAT'S FOR BREAKFAST?

Breakfast eaters experience lower blood pressures than breakfast skippers.²⁷⁹ In planning breakfast, be warned of the "cereal killer". In laboratory animals, most processed breakfast cereals, dry or hot, have a hard time sustaining life and cause hypertension.²⁸⁰ Choose whole plant based unrefined foods for a healthy, blood pressure friendly breakfast. If you cannot, by looking at the breakfast food, determine its identity, such as "O"s or flakes which do not resemble their origin, don't put it in your mouth. It is much better to eat identifiable cereals such as granola, oatmeal, and other whole grain cereal foods.

The reason why refining foods causes so much hypertension is that fiber and minerals are removed in the process.²⁸¹ Fiber from grains has been shown to reduce the risk of hypertension by 40%.²⁸² Vegetable fiber, 65 gm/day, can reduce your blood pressure by 12-14 points.²⁸³

Beans help lower blood pressure because they are low in sodium, have healthy fiber and contain the blood pressure lowering minerals potassium, magnesium, and calcium.²⁸⁴

Even nuts are beneficial. A daily serving of nuts can reduce the risk of hypertension by 18%.²⁸⁵ The preferred variety is raw with little or no salt. Even

“soy nuts”, a roasted soybean product, have been shown to positively impact blood pressure.²⁸⁶

One important antioxidant vitamin for addressing blood pressure is vitamin C. When blood levels are maintained from natural dietary sources, blood pressure drops,²⁸⁷ but not when blood levels are attempted through artificial (pill supplementation) sources.²⁸⁸ Diets high in this and other naturally occurring vitamins from fruit and vegetables lower blood pressure.²⁸⁹ One such diet is the Hawaiian diet. This diet is high in complex carbohydrate (77% of calories), low in fat (12% of calories), moderate in protein (11% of calories), and has been shown to have decreases in blood pressure by- as much as 10 points.²⁹⁰

One research group, wanting to maximize dietary impact on blood pressure management, put their patients on a six-month 62% uncooked fresh fruit and vegetable diet. The result was an 8-pound weight loss and an 18 mm Hg blood pressure decrease. In passing, the researchers noted that 80% of those in this study who smoked or drank alcohol abstained spontaneously. They reported increased sensitivity to alcohol, cigarette smoke, chemical odors, and medications. Several also had nausea, vomiting, and malaise after typical restaurant or banquet dinners.²⁹¹ What kind of diet are we talking about? The original diet! “And God said, Behold, I have given you every herb bearing seed, which is upon the face of all the earth, and every tree, in the which is the fruit of a tree yielding seed; to you it shall be for meat.” “and thou shalt eat the herb of the field;”²⁹² “Grains, fruits, nuts, and vegetables constitute the diet chosen for us by our Creator. These foods, prepared in as simple and natural a manner as possible, are the most healthful and nourishing. They impart a strength, a power of endurance, and a vigor of intellect that are not afforded by a more complex and stimulating diet.”²⁹³

WHY EAT SO MUCH?

“And put a knife to thy throat, if thou be a man given to appetite.”²⁹⁴ Hypertension increases with overeating.^{295,296} Do not overeat! “...eat in due season, for strength, and not for drunkenness!”²⁹⁷ Eating less food (caloric restriction) reduces blood vessel stiffness, improves vascular relaxation and lowers blood pressure.^{298,299}

Taking it a step farther, fasting has been shown to be an effective modality is lowering blood pressure. You can jump start your blood pressure reduction with a water only fast.³⁰⁰ Fasting effectively reduces stubborn hypertension.³⁰¹

Fasting one or two days a week may be more effective than pills.³⁰²

HAVE SOME LEMON IN YOUR WATER!

With dehydration as one of the causes of hypertension,³⁰³ what is the best way to get my water? Adding fresh squeezed lemon to water maximizes its impact on hypertension.³⁰⁴ We recommend that you squeeze the juice of one lemon into your first quart of water for the day and drink it at least 30 minutes before breakfast to lower blood pressure. We recommend that the water be lukewarm, not hot or cold.

TACHYCARDIA: INCREASED HEART RATE

If the heart beats faster, a higher volume of blood is pumped and the blood pressure rises.³⁰⁵ We call this tachycardia. Anything that causes the resting heart rate to increase accelerates the risk of hypertension. In fact, for every 10 beats/min increase in heart rate the risk of hypertension increases by 42%.³⁰⁶ You see, athletes have very low heart rates. People who have not been exercising have a high heart rate, and a high risk of hypertension. Stress also raises the pulse rate.

DOES STRESS RUN IN YOUR BLOOD?

Are you easily startled? This is part of your response to stress and is an early sign you may be headed for hypertension.³⁰⁷ Besides raising the heart rate and tightening the blood vessels, another way stress causes hypertension and blood clots is that it thickens the blood so that it requires more pressure to pump it through the blood vessels.³⁰⁸ People who never get anxious have a significantly lower incidence of hypertension.³⁰⁹ “Take therefore no thought for the morrow: for the morrow shall take thought for the things of itself. Sufficient unto the day is the evil thereof.”³¹⁰

Another way to look at it is that people who handle stressful life events more effectively have greater success maintaining healthy blood pressure.³¹¹ This may have something to do with personality. The easy-going type B personalities go easier on blood pressure than their type A counterparts.³¹² Easy-going, laid-back, calm, relaxed responses to life’s challenges predict normal blood pressure.³¹³ People with cool tempers experience calmer blood pressures³¹⁴ and fewer heart attacks.³¹⁵ “He that is slow to anger is better than the mighty; and he that ruleth his spirit than he that taketh a city.”³¹⁶ What’s more, happier more cheerful people have lower blood pressures.³¹⁷

Having trouble coping? Stress management training, including hostility reduction and anger management have been shown to be effective at lowering blood pressure.³¹⁸

Does keeping an animal help blood pressure? Yes, loving pets have a blood pressure lowering effect for people with hypertension.

Of course, stress is how you perceive your risks, not the risks themselves. People who experience events as negative have a higher risk of hypertension.³¹⁹ The question then is; how can I avoid, or get over stress? How do I change my perceptions of risk away from the negative toward the positive? The first step is being aware when stress is playing a role in your life. Does your pulse rise? Are you physically tense? Do you experience headaches or anxiety? Are you nervous? Does fatigue dog your steps? Do you have high blood pressure? Being in touch with your emotions is key because stress is an emotional experience.

When you discover that you have stress, your next step is to realize or remember what thoughts surround or accompany your stress. Then, evaluate the validity of those thoughts. Are they rational? Are they negative or are they positive? Once you have flushed out the thoughts underlying your stressful feelings the next step is to determine what these thoughts tell you about what you believe. Beliefs underlie thoughts, thoughts underlie feelings and feelings drive actions or behaviors like hypertension. Take a closer look at your secretly held beliefs, especially the ones underlying stressful feelings. Most often you will discover beliefs that set you up for unreasonable fear, worry, or anger. You need to challenge these beliefs and replace them with rational ones. "And ye shall know the truth, and the truth shall make you free."³²⁰

Beliefs that you will find most helpful for achieving good health are ones obtained from the Bible where we learn that "There is no fear in love; but perfect love casteth out fear: because fear hath torment. He that feareth is not made perfect in love."³²¹

One common stress provoking fear is one of financial failure. But when we have turned our lives over to God, we can expect His care. "And he said unto his disciples, Therefore I say unto you, Take no thought for your life, what ye shall eat; neither for the body, what ye shall put on. The life is more than meat, and the body is more than raiment. Consider the ravens: for they neither sow nor reap; which neither have storehouse nor barn; and God feedeth them: how much more are ye better than the fowls?

And which of you with taking thought can add to his stature one cubit? If ye then be not able to do that thing which is least, why take ye thought for the rest? Consider the lilies how they grow: they toil not, they spin not; and yet I say unto you, that Solomon in all his glory was not arrayed like one of these. If then God so clothe the grass, which is to day in the field, and tomorrow is cast into the oven; how much more will he clothe you, O ye of little faith? And seek not ye what ye shall eat, or what ye shall drink, neither be ye of doubtful mind. For all these things do the nations of the world seek after and your Father knoweth that ye have need of these things. But rather, seek ye the kingdom of God; and all these things shall be added unto you."³²²

People who put the past behind them, not ruminating about past anger-provoking events are at lower risk for hypertension and its complications.³²³ "Brethren, I count not myself to have apprehended: but this one thing I do, forgetting those things which are behind, and reaching forth unto those things which are before, I press toward the mark for the prize of the high calling of God in Christ Jesus."³²⁴

STRESSFUL LIFE EVENTS

One way to increase your stress is to subject your mind to the impressions being made on television. Two (2) hours a day of television watching increases the risk of hypertension by 40%.³²⁵ As I am sure you are well aware, if you think about it, the main character of the plot is either in trouble, getting into trouble, or getting out of trouble, and life just is not that bad.

Difficulty paying medical expenses is associated with increased hypertension.³²⁶ It is also the number one reason for bankruptcy in the United States.³²⁷

WORK PLEASURE

The rewards of gainful, meaningful employment bear mentioning. People happy with their jobs and their income are more likely to experience happy, healthy blood pressure.^{328,329} People who enjoy their jobs and are not overworked or depressed have lower blood pressures.³³⁰ Having purpose in life helps moderate blood pressure.³³¹

SOCIAL PLEASURE

Looking at the social aspects of blood pressure control, people with close supportive friends have lower blood pressures.³³² Indeed, sharing a negative life experience with an ambivalent friend raises blood pressure and heart rate. Alternatively sharing

the same event with a caring, supportive friend lowers blood pressure and heart rate. Similarly, blood pressure levels are lowest when people are with family and highest when among strangers.³³³

PET POWER

Does keeping an animal help blood pressure? Yes, loving pets have a blood pressure lowering effect for people with hypertension.³³⁴

MUSIC MAGIC

Do you enjoy listening to music? Music can help blood pressure.³³⁵ Blood pressures respond positively to classical music, but not jazz or pop.³³⁶

EMERGING URBAN DANGER

Environment plays a significant role in stress. Moving to the city? Expect a 23-point rise in your systolic blood pressure and a 9-point rise in your diastolic blood pressure.³³⁷ There are hazards in just commuting to a city. Traffic related air pollution and noise significantly increase the risk of hypertension.³³⁸ People living in small rural towns have half the risk of hypertension as people living in large, industrialized cities.³³⁹

RURAL RELIEF

Quietness, solitude and silence soothe the nerves and lower your blood pressure.^{340,341} The stillness is remedial, "Be still, and know that I am God."³⁴² People living where the din of busy roads never reaches their ears have a 37% lower risk of hypertension.³⁴³ Outdoor walks, enjoying trees, flowers, and other wonders of God's great nature, lower blood pressure and stress.³⁴⁴ Even bringing the outdoors into the home through houseplants has been shown to improve blood pressure.^{345,346}

LIFE'S PLEASURE: HELPING OTHERS

If you have extra time and want to do something beneficial for your blood pressure, volunteering is a wonderful aid in keeping blood pressures normal.³⁴⁷ Find someone else who could use your help and give of yourself to the needs of the world. "It is more blessed to give than to receive."³⁴⁸

FREEDOM FROM GUILT AND RESENTMENT

Guilt and resentment cause hypertension. Freedom from guilt³⁴⁹ and resentment³⁵⁰ lowers blood pressure. Okay, how do I do that? Easier said

than done! Well, it is not really something you can do on your own. You are going to need help. Only God can accomplish this, with your cooperation. Are you ready?

Guilt comes from sin. "Sin is the transgression of the law."³⁵¹ It is being out of harmony with the loving God of nature and the universe, of this you must first be conscious. You may already have a sense that something is not right, that you could use some more peace in your life.

How do I know if I have guilt? One of two ways: 1. Compare your life to the standard of God's holy law³⁵² and see if there is a disparity between what you are and what it enjoins, "for by the law is the knowledge of sin."³⁵³ Go to the law, read the first commandment, and ask God to reveal to you if you are in harmony with it. Then proceed to the following commandments and ask the same question. 2. Compare your life with that of Jesus Christ, "Who did no sin."³⁵⁴ meaning that His life was an example of what it is like to be at complete peace with God and keep His law free of guilt. Read His biographies at the beginning of the New Testament (Mathew, Mark, Luke, John) of the Bible and ask God to reveal to you the differences between what you are studying and what your life could have been like if you had been like Jesus. This is another way of comparing your life to what the law enjoins and discovering where you may have sinned.

If, after trying this, you find yourself believing that you are problem free, it may only indicate that you have undertaken this comparison too superficially and need to spend a little more time with it. A thorough investigation will reveal that, "all have sinned, and come short of the glory of God."³⁵⁵ And, "If we say that we have no sin, we deceive ourselves, and the truth is not in us."³⁵⁶

God's Holy Spirit will help you and your conscience will be awakened to see the evil of sin, its power and guilt. Sin separates you from God and peace and brings you into bondage. It brings feelings of guilt that result in physical symptoms like hypertension. The more you struggle on your own to escape, "the more you realize your helplessness. Your motives are impure; your heart is unclean. You see that your life has been filled with selfishness and sin. You long to be forgiven, to be cleansed, and to be set free. Harmony with God, likeness to Him – what can you do to obtain it?"³⁵⁷

"It is peace that you need – Heaven's forgiveness, peace, and love in the soul. Money cannot buy it, intellect cannot procure it, wisdom cannot attain to it; you can never hope, by your own efforts to secure it. But God offers it to you as a gift, "without money and without price." Isaiah 55:1. It is yours if you will

but reach out your hand and grasp it. The Lord says, "Though your sins be as scarlet, they shall be as white as snow; though they be red like crimson, they shall be as wool." Isaiah 1:18. "A new heart also will I give you, and a new spirit will I put within you. Ezekiel 36:26."³⁵⁷

Guilt and resentment cause hypertension. Freedom from guilt and resentment lowers blood pressure.

Confess your sins, and in heart put them away. Say, "Dear God, I have sinned and have suffered the results of a sense of guilt and have the disease of high blood pressure. Please forgive me and give me a new heart to know you and live free from offence toward you and others." Resolve to give yourself to God. "Now go to Him and ask that He will wash away your sins and give you a new heart. Then believe that He does this because He has promised. This is the lesson which Jesus taught while He was on earth, that the gift which God promises us, we must believe we do receive, and it is ours. Jesus healed the people of their diseases when they had faith in His power; He helped them in the things which they could see, thus inspiring them with confidence in Him concerning things which they could not see--leading them to believe in His power to forgive sins. This He plainly stated in the healing of the man sick with palsy: 'That ye may know that the Son of man hath power on earth to forgive sins, (then saith He to the sick of the palsy,) Arise, take up thy bed, and go unto thine house.' Matthew 9:6. So also John the evangelist says, speaking of the miracles of Christ, 'These are written, that ye might believe that Jesus is the Christ, the Son of God; and that believing ye might have life through His name.' John 20:31."³⁶⁰

To maintain peace and freedom from blood-pressure-raising guilt, "Fight the good fight of faith,"³⁵⁸ believe that God has taken care of your past sins, daily read His word to learn new things that will help keep blood-pressure-raising guilt away, and "press toward the mark for the prize of the high calling of God in Christ Jesus."³⁵⁹ "To them who by patient continuance in well doing seek for glory and honor, and immortality, eternal life:"³⁶⁰ Rest in the assurance of eternal life and a good afterlife.

Resentment: resentment arises out of misunderstanding between people. It comes from not forgiving someone else. It is a form of anger. Resentment grows if not checked. Do not allow the wound to fester and break out in poisoned words, which taint the minds of those who hear. Do not allow bitter thoughts to continue to fill your mind.

"Thou shalt not avenge, nor bear any grudge against the children of thy people, but thou shalt love thy neighbour as thyself: I am the LORD."³⁶¹

Go to your brother, and in humility and sincerity talk with him about the matter. "Moreover, if thy brother shall trespass against thee, go and tell him his fault between thee and him alone: if he shall hear thee, thou hast gained thy brother. But if he will not hear thee, then take with thee one or two more, that in the mouth of two or three witnesses every word may be established. And if he shall neglect to hear them, tell it unto the church: but if he neglect to hear the church, let him be unto thee as an heathen man and a publican."³⁶² At this point you have done your part, no matter what the outcome, choose to let God have the feelings of resentment and bitterness, and fill your mind with glad saying like from the Bible books of Psalms or Proverbs.

PATIENCE AND FORGIVENESS LOWER BLOOD PRESSURE

People who cultivate the characteristic of patience enjoy much more normal blood pressures.³⁶³ This reminds me of a very important passage in the last book of the Bible, "Here is the patience of the saints: here are they that keep the commandments of God, and the faith of Jesus."³⁶⁴

A spirit of forgiveness has been shown to bring blood pressure down.³⁶⁵ "Forgiving one another, even as God for Christ's sake hath forgiven you."³⁶⁶

RELIGION, BIBLE STUDY, AND PRAYER LOWER BLOOD PRESSURE

In stressful situations, prayer has been shown to lower blood pressure.³⁶⁷ Further, people who attend religious services,³⁶⁸ and pray or study the Bible frequently, have a 40% lower risk of hypertension.³⁶⁹ People who substituted religious media (TV or Radio) for personal experience and social contact with other believers suffered elevated blood pressures.³⁷²

Most people find wars and natural disasters like hurricanes and earthquakes stressful. In the wake of earthquakes and other stress-raising natural disasters more people come down with high blood pressure.³⁷⁰ I would advise you to avoid this source of stress, but I would be misleading you if I asserted that these situations were going to become less frequent. "For nation shall rise against nation, and kingdom against kingdom: and there shall be earthquakes in diverse places, and there shall be famines and troubles: these are the beginnings of sorrows."³⁷¹ In other words, you have not seen

anything yet. And as we get closer to the second coming of Jesus Christ these things will become more and more common. Stress management needs to be God derived to meet such impending devastation.

DOES LOVE LAST FOREVER?

We have already discussed the impact of major stressful life events on the development of high blood pressure. One such major stressful life event is the loss of a loved one. A study out of the University of Michigan, Ann Arbor, revealed that older adults who lost a loved one to death, but who believe in a good afterlife were much less likely to develop hypertension.³⁷² This raises an important question; do we have reason to believe in a good afterlife? If not, we are all headed for high blood pressure. If God is just waiting, as many religions teach, to roast sinners in the fires of hell for eternity, stage II hypertension here we come! But what is the truth about the afterlife?

Jesus was right up front in telling us, that after He left this earth, He would be preparing to have us join Him. "Let not your heart be troubled: ye believe in God, believe also in me. In my Father's house are many mansions: if it were not so, I would have told you. I go to prepare a place for you. And if I go and prepare a place for you, I will come again, and receive you unto myself; that where I am, there ye may be also."³⁷³ When are we going to join Him? When do we get this reward? "For the Son of man shall come in the glory of his Father with his angels; and then he shall reward every man according to his works."³⁷⁴ He will be rewarding everyone at His second coming.

How will He know who to give the good rewards to and who to exclude? "I charge thee therefore before God, and the Lord Jesus Christ, who shall judge the quick and the dead at his appearing and his kingdom;" "Henceforth there is laid up for me a crown of righteousness, which the Lord, the righteous judge, shall give me at that day: and not to me only, but unto all them also that love his appearing."³⁷⁵ God makes a decision on that in a court setting called the judgment.³⁷⁶ Clearly no judgment has been made or reward given at this point in history, it does not occur until His second coming. Even Paul waits for that day to receive his "crown of righteousness". So, all, dead or alive, good or evil, are still waiting for the judgment and their reward.

What about those who have already died? Jesus said, "Marvel not at this: for the hour is coming, in the which all that are in the graves shall hear his voice, And shall come forth; they that have done

good, unto the resurrection of life; and they that have done evil, unto the resurrection of damnation."³⁷⁷ Jesus has yet to raise the dead so that they can receive their reward. No one is currently burning in hell or enjoying heaven.

What are the dead doing right now? Can I talk to one of them? Will any of them remember me? What says the Scriptures concerning these things? The beloved David declares that man is not conscious in death. "His breath goeth forth, he returneth to his earth; in that very day his thoughts perish."³⁷⁸ "In death there is no remembrance of thee; in the grave who shall give thee thanks?" "The dead praise not the Lord, neither any that go down into silence."³⁷⁹ Solomon bears the same testimony: "The living know that they shall die; but the dead know not anything." "Their love, and their hatred, and their envy, is now perished; neither have they any more a portion forever in anything that is done under the sun." "There is no work, nor device, nor knowledge, nor wisdom, in the grave, whither thou goest."³⁸⁰

When will this resurrection take place? "For this we say unto you by the word of the Lord, that we which are alive (and) remain unto the coming of the Lord shall not prevent them which are asleep. For the Lord himself shall descend from heaven with a shout, with the voice of the archangel, and with the trump of God: and the dead in Christ shall rise first: Then we which are alive (and) remain shall be caught up together with them in the clouds, to meet the Lord in the air: and so shall we ever be with the Lord. Wherefore comfort one another with these words."³⁸¹ That is really comforting and helps produce good blood pressure.

Even the Old Testament character Job was aware of this future resurrection: "If a man die, shall he live again? All the days of my appointed time will I wait, till my change come. Thou shalt call, and I will answer thee: thou wilt have a desire to the work of thine hands."³⁸²

What change was Job anticipating at the resurrection? Paul answers this for us, "In a moment, in the twinkling of an eye, at the last trump: for the trumpet shall sound, and the dead shall be raised incorruptible, and we shall be changed. For this corruptible must put on incorruption, and this mortal must put on immortality. So, when this corruptible shall have put on incorruption, and this mortal shall have put on immortality, then shall be brought to pass the saying that is written, Death is swallowed up in victory. O death, where is thy sting? O grave, where is thy victory? The sting of death is sin; and the strength of sin is the law. But thanks be to God, which giveth us the victory through our Lord Jesus Christ. Therefore, my beloved brethren, be ye steadfast, unmoveable,

always abounding in the work of the Lord, forasmuch as ye know that your labor is not in vain in the Lord."³⁸³ If you are faithful to God, you have the promise of a new body at Jesus' return, free from diseases such as hypertension.

The outcome of the judgment at the second coming of Jesus is that the righteous go to heaven and the wicked to hell. For some people the resurrection is the beginning of a happy afterlife, these are the ones we discussed earlier who believe in a good afterlife and experience lower blood pressure now. "And many of them that sleep in the dust of the earth shall awake, some to everlasting life, and some to shame and everlasting contempt."³⁸⁴

If you are faithful to God, you have the promise of a new body at Jesus' return, free from diseases such as hypertension.

And what is the punishment of those who choose to reject God's offer of forgiveness and mercy? Is hell forever? Contemplating hell is daunting, and the thought of being condemned to go there could certainly raise blood pressure. Why should there even be a hell? "Then shall he say also unto them on the left hand, depart from me, ye cursed, into everlasting fire, prepared for the devil and his angels."³⁸⁵ Hell was never intended for you or me. It is for the devil and his angels who oppose God. But, if we join the devil in sin, in fighting the righteous law of God, then we are choosing to join the devil in his plight. Really, God would rather we repent and choose His way of life: "The Lord is...longsuffering to us-ward, not willing that any should perish, but that all should come to repentance."³⁸⁶

Do the wicked suffer in the fires of hell through all eternity? "For, behold, the day cometh, that shall burn as an oven; and all the proud, yea, and all that do wickedly, shall be stubble: and the day that cometh shall burn them up, saith the LORD of hosts, that it shall leave them neither root nor branch. And ye shall tread down the wicked; for they shall be ashes under the soles of your feet in the day that I shall do this, saith the LORD of hosts."³⁸⁷ So really, although it is a fearful ordeal, it is self-limited, it comes to an end, all that is left is ashes. Matthew speaks of this, "Whose fan is in his hand, and he will thoroughly purge his floor, and gather his wheat into the garner; but he will burn up the chaff with unquenchable fire."³⁸⁸ Burned Up!

Has this ever happened before? It has: "Even as Sodom and Gomorrah, and the cities about them in like manner, giving themselves over to fornication, and going after strange flesh, are set forth for an example, suffering the vengeance of eternal fire."³⁸⁹

These cities were very rebellious against God and His Law so God destroyed them with eternal fire as an example of hell. Are they still burning? Of course not, but they did burn to ashes, to be trodden under foot, as the wicked will be.

Okay, but what if my loved ones choose a life which leads them to hell, won't this be on my mind through all eternity? No, "And God shall wipe away all tears from their eyes"³⁹⁰ "For, behold, I create new heavens and a new earth: and the former shall not be remembered, nor come into mind."³⁹¹

Will God bring a complete end to all sorrow, pain, the high blood pressure? "What do ye imagine against the LORD? he will make an utter end: affliction shall not rise up the second time."³⁹²

While those who do not embrace God's ways and law are "as though they had not been."³⁹³, the righteous will be enjoy being with Jesus and eating from the Tree of Life whose leaves are for healing, "In the midst of the street of it, and on either side of the river, was there the tree of life, which bare twelve manner of fruits, and yielded her fruit every month: and the leaves of the tree were for the healing of the nations."³⁹⁴ Perhaps for healing even of hypertension. We know that there will be no disease in heaven. "And God shall wipe away all tears from their eyes; and there shall be no more death, neither sorrow, nor crying, neither shall there be any more pain: for the former things are passed away."³⁹⁵ What a wonderful day that will be! Don't you look forward to that—Total freedom from death, sorrow, crying and pain? Are you ready to be healed?

REVIEW AND RECAP

Review: Blood pressure is a product of the pumping of the heart, the size of the blood vessel through which the blood must flow, the thickness of the blood, and the volume of blood to be pumped.

A fast heart rate pumps more blood than normal and increases blood pressure. We call this tachycardia. Common causes of tachycardia are stress, having poor cardiovascular fitness (being out of shape), caffeine, tobacco and dehydration.

If the blood vessels tighten up or constrict, making the blood vessel size smaller, higher blood pressure is needed to get the required amount of blood to its destination. We refer to this tightening up effect as vasoconstriction. Vasoconstriction is caused by psychological and physical stress, thermal stress from cold-exposure, caffeine, tobacco, and dehydration.

Blood vessels normally expand and relax with each heartbeat. If the blood vessels become hard, their stiffness resists the free flow of blood and up

goes the blood pressure. Atherosclerosis is an example of this process, so are sugar glycation, endothelial dysfunction, and stiffness of the blood vessels running through muscles from a sedentary lifestyle.

If the blood becomes thick and sludgy, more pressure is required to carry it through the blood vessels and hypertension results. When blood gets thick, we say that the viscosity has increased too much. Examples of the blood-thickening threats include dehydration, overeating, psychological stress, a diet too rich in fat and refined carbohydrates, and high blood cholesterol or triglycerides.

If the blood vessels are being choked by something pressing on them from their sides, the effect is like putting your thumb over the end of a garden hose, the result is increased blood pressure. We call this external compression. Things that press on the blood vessels include tissue swelling, atherosclerotic plaque, sugar coating called glycation (usually from diabetes), inflammation, and especially tight clothing like belts and elastic bands.

Finally, if the volume of blood increases, this increases the amount of blood entering the heart, which in turn increases the amount of blood leaving the heart and with each beat this increases the overall blood pressure. We often refer to this phenomenon as fluid retention. Lifestyle habits causing volume overload include eating too much salt, obesity, a sedentary lifestyle, poorly clad chilled extremities, and kidney failure.

SUMMARY: LETTING HYPERTENSION GO!

- Keep well hydrated with pure vitalizing water; add a little fresh lemon juice, too.
- Take regular time for enjoyable exercise in the great outdoors, far from traffic, pollution, cities and stressful crowds.
- Get plenty of warm sunshine and invigorating fresh air.
- Eat a delicious unrefined whole-plant-based diet; high in fiber, minerals, vitamins, and antioxidants, low in salt, with no refined carbohydrates, processed fats, or hypertension-causing animal products.
- Avoid the pitfalls of tobacco, caffeine, alcohol, drugs, fried foods, fructose, sodium, MSG, and overeating.
- Make weight control a habit.
- Guard your sleep and evaluate its quality for effectiveness.
- Let God know you need Him to release you from stress, guilt, fear, and hypertension. Rely on Him for guidance as you come more into line with His original plan for diet and lifestyle.

For further ideas on how to incorporate what you have just learned into your daily life, see the article entitled, "How Can I Apply Healthy Principles in My Daily Life", or Lifestyle Choices.

¹ Suhr JA, Patterson SM, Austin AW, Heffner KL. The relation of hydration status to declarative memory and working memory in older adults. *J Nutr Health Aging*. 2010 Oct;14(10):840-3.

² Khokhar AM, Slater JD. Increased renal excretion of arginine-vasopressin during mild hyponatremia in young men with mild essential benign hypertension. *Clin Sci Mol Med Suppl*. 1976 Dec;3:691s-694s.

³ Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, Jones DW, Materson BJ, Oparil S, Wright JT Jr, Rocella EJ, National Heart, Lung, and Blood Institute Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure; National High Blood Pressure Education Program Coordinating Committee. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA*. 2003 May 21;289(19):2560-72.

⁴ Thornton SN. Thirst and hydration: physiology and consequences of dysfunction. *Physiol Behav*. 2010 Apr 26;100(1):15-21.

⁵ Roudaut R, Gosse P, Aouizerate E, Dalocchio M. Low blood pressure. *Ann Cardiol Angeiol (Paris)*. 1989 May;38(5):279-80.

⁶ Pearce KA, Furberg CD, Rushing J. Does antihypertensive treatment of the elderly prevent cardiovascular events or prolong life? A meta-analysis of hypertension treatment trials. *Arch Fam Med*. 1995 Nov;4(11):943-9.

⁷ Franco OH, Peeters A, Bonneux L, de Laet C. Blood pressure in adulthood and life expectancy with cardiovascular disease in men and women: life course analysis. *Hypertension*. 2005 Aug;46(2):280-6. Epub 2005 Jun 27.

⁸ van Dijk EJ, Breteler MM, Schmidt R, Berger K, Nilsson LG, Oudkerk M, Pajak A, Sans S, de Ridder M, Dufouil C, Fuhrer R, Giampaoli S, Launer LJ, Hofman A; CASCADE Consortium. The association between blood pressure, hypertension, and cerebral white matter lesions: cardiovascular determinants of dementia study. *Hypertension*. 2004 Nov;44(5):625-30.

⁹ van Swieten JC, Geyskes GG, Derix MM, Peck BM, Ramos LM, van Latum JC, van Gijn J. Hypertension in the elderly is associated with white matter lesions and cognitive decline. *Ann Neurol*. 1991 Dec;30(6):825-30.

¹⁰ Longstreth WT Jr, Arnold AM, Beauchamp NJ Jr, Manolio TA, Lefkowitz D, Jungreis C, Hirsch CH, O'Leary DH, Furberg CD. Incidence, manifestations, and predictors of worsening white matter on serial cranial magnetic resonance imaging in the elderly: the Cardiovascular Health Study. *Stroke*. 2005 Jan;36(1):56-61.

¹¹ Wilburn AJ, King DS, Glisson J, Rockhold RW, Wofford MR. The natural treatment of hypertension. *J Clin Hypertens (Greenwich)*. 2004 May;6(5):242-8.

¹² Sivaprakasapillai B, Edirisinghe I, Randolph J, Steinberg F, Kappagoda T. Effect of grape seed extract on blood pressure in subjects with the metabolic syndrome. *Metabolism*. 2009 Dec;58(12):1743-6.

¹³ Häckel LP, Cuttler G, Dovich SS, Lima-Landman MT, Nicolau M. Inhibition of angiotensin-converting enzyme by quercetin alters the vascular response to bradykinin and angiotensin I. *Pharmacology*. 2002 Aug;65(4):182-6.

¹⁴ Rosenfeld FL, Haas SJ, Krum H, Hadji A, Ng K, Leong JY, Watts GF. Coenzyme Q10 in the treatment of hypertension: a meta-analysis of the clinical trials. *J Hum Hypertens*. 2007 Apr;21(4):297-306.

¹⁵ Kwon YI, Vatter DM, Shetty K. Evaluation of clonal herbs of Lamiales species for management of diabetes and hypertension. *Asia Pac J Clin Nutr*. 2006;15(1):107-18.

¹⁶ Apostolides E, Kwon YI, Shetty K. Potential of cranberry-based herbal synergies for diabetes and hypertension management. *Asia Pac J Clin Nutr*. 2006;15(3):433-41.

¹⁷ El Bardai S, Lyoussi B, Wilbo M, Morel N. Pharmacological evidence of hypotensive activity of *Marrubium vulgare* and *Foeniculum vulgare* (Fennel) in spontaneously hypertensive rats. *Clin Exp Hypertens*. 2001 May;23(4):329-43.

¹⁸ Jadhav RB, Bhatnagar SP, Surana SJ. Diuretic activity of squamate mistletoe, *Viscum angulatum*. *Pharm Biol*. 2010 Apr;48(4):417-21.

¹⁹ Ye F, Du ZG, Cui AQ, Lu XT. Study on the mechanism of compound mistletoe fluidextract in relieving hypertension. *J Tradit Chin Med*. 2009 Dec;29(4):291-5.

²⁰ Imenshahidi M, Hosseinzadeh H, Javadpour Y. Hypotensive effect of aqueous saffron extract (*Crocus sativus* L.) and its constituents, safranal and crocin, in normotensive and hypertensive rats. *Phytother Res*. 2010 Jul;24(7):990-4.

²¹ Walker AF, Marakis G, Simpson E, Hope JL, Robinson PA, Hassanein M, Simpson HC. Hypotensive effects of Hawthorn for patients with diabetes taking prescription drugs: a randomised controlled trial. *Br J Gen Pract*. 2006 Jun;56(527):437-43.

²² Circostra C, De Pasquale R, Samperi S, Pino A, Occhiuto F. Biological and analytical characterization of two extracts from *Valeriana officinalis* (Valerian). *J Ethnopharmacol*. 2007 Jun 13;112(2):361-7.

²³ Katarinen MJ, Puska PM, Korhonen MH, Mustonen JN, Salomaa VV, Sundvall JE, Tuomilehto JO, Uusitupa MJ, Nissinen AM, LIFE Study Group. Non-pharmacological treatment of hypertension in primary health care: a 2-year open randomized controlled trial of lifestyle intervention against hypertension in eastern Finland. *J Hypertens*. 2002 Dec;20(12):2505-12.

²⁴ García-Ruiz PJ, Javier Jiménez-Jiménez F, García de Yébenes J. Calcium channel blocker-induced parkinsonism: clinical features and comparisons with Parkinson's disease. *Parkinsonism Relat Disb*. 1998 Dec;4(4):211-214.

²⁵ Skoog I, Lernfelt B, Landahl S, Palmertz B, Andreasson LA, Nilsson L, Persson G, Odén A, Svanborg A. 15-year longitudinal study of blood pressure and dementia. *Lancet*. 1996 Apr 27;347(9009):1141-5.

²⁶ Longstreth WT Jr, Arnold AM, Beauchamp NJ Jr, Manolio TA, Lefkowitz D, Jungreis C, Hirsch CH, O'Leary DH, Furberg CD. Incidence, manifestations, and predictors of worsening white matter on serial cranial magnetic resonance imaging in the elderly: the Cardiovascular Health Study. *Stroke*. 2005 Jan;36(1):56-61.

²⁷ Salerno JA, Murphy DG, Horwitz B, DeCarli C, Haxby JV, Rapoport SI, Schapiro MB. Brain atrophy in hypertension. A volumetric magnetic resonance imaging study. *Hypertension*. 1992 Sep;20(3):340-8.

²⁸ Lubianca JN, Faccin CS, Fuchs FD. Oral contraceptives: a risk factor for uncontrolled blood pressure among hypertensive women. *Contraception*. 2003 Jan;67(1):19-24.

²⁹ Lubianca JN, Moreira LB, Gus M, Fuchs FD. Stopping oral contraceptives: an effective blood pressure-lowering intervention in women with hypertension. *J Hum Hypertens*. 2005 Jun;19(6):451-5.

³⁰ Licht CM, de Geus EJ, Seldenrijk A, van Hout HP, Zitman FG, van Dyck R, Penninx BW. Depression is associated with decreased blood pressure, but antidepressant use increases the risk for hypertension. *Hypertension*. 2009 Apr;53(4):631-8.

³¹ Forman JP, Rimm EB, Curhan GC. Frequency of analgesic use and risk of hypertension among men. *Arch Intern Med*. 2007 Feb 26;167(4):394-9.

³² Laine L, White WB, Rostom A, Hochberg M. COX-2 selective inhibitors in the treatment of osteoarthritis. *Semin Arthritis Rheum*. 2008 Dec;38(3):165-87.

³³ Gaziano JM. Nonnarcotic analgesics and hypertension. *Am J Cardiol*. 2006 May 8;97(9A):10-6. Epub 2006 Mar 30.

³⁴ Forman JP, Stampfer MJ, Curhan GC. Non-narcotic analgesic dose and risk of incident hypertension in US women. *Hypertension*. 2005 Sep;46(3):500-7.

³⁵ Sudano I, Flammer AJ, Périer D, Enseleit F, Hermann M, Wolfram M, Hirt A, Kaiser P, Hurlimann D, Neidhart M, Gay S, Holzmeister J, Nussberger J, Mocharfa P, Landmesser U, Haile SR, Corti R, Vanhoute PM, Lüscher TF, Noll G, Ruschitzka F. Acetaminophen increases blood pressure in patients with coronary artery disease. *Circulation*. 2010 Nov 2;122(18):1789-96.

³⁶ Osthege Y, Yoon SS, Hughes J, Louis T. Hypertension awareness, treatment, and control - continued disparities in adults: United States, 2005-2006. NCHS data brief no Hyattsville, MD: National Center for Health Statistics. 2008.

³⁷ Nedley N. Proof Positive: How to Reliably Combat Disease and Achieve Optimal Health through Nutrition and Lifestyle (Ardmore, OK: Nedley Publishing, 1999).

³⁸ Okken VS, Niemeijer MG, Dijkstra A, Baars MW, Said S, Hoogenberg K, Orfgen H, Otten S, Cleophas TJ. The effect of physical, social and psychological factors on drug compliance in patients with mild hypertension. *Neth Heart J*. 2008 Jun;16(6):297-300.

³⁹ Haidler AW, Larson MG, Franklin SS, Levy D. Systolic blood pressure, diastolic blood pressure, and pulse pressure as predictors of risk for congestive heart failure in the Framingham Heart Study. *Ann Intern Med*. 2003 Jan 7;138(1):10-6.

⁴⁰ Kannel WB, Schwartz MJ, McNamara PM. Blood pressure and risk of coronary heart disease: the Framingham Study. 1969. *Chest*. 2009 Nov;136(5 Suppl):e23.

⁴¹ Law M, Wald N, Morris J. Lowering blood pressure to prevent myocardial infarction and stroke: a new preventive strategy. *Health Technol Assess*. 2003;7(31):1-94.

⁴² Conen D, Ridker PM, Buring JE, Glynn RJ. Risk of cardiovascular events among women with high normal blood pressure or blood pressure progression: prospective cohort study. *BMJ*. 2007 Sep 1;335(7617):432.

⁴³ Hiramoto JS, Howell B, Reilly LM, Chuter TA. Effect of systemic blood pressure on aneurysm size in the presence of a type II endoleak. *Vascular*. 2008 Nov-Dec;16(6):321-5.

- 44 Neal B, MacMahon S, Chapman N: Effects of ACE inhibitors, calcium antagonists, and other blood-pressure-lowering drugs: results of prospectively designed overviews of randomised trials. *Blood Pressure Lowering Treatment Trialists' Collaboration*. *Lancet*. 2000 Dec 9;356(9246):1955-64.
- 45 Strano A, Novio S, Avellone G, Di Garbo V, Abrignani MG, Liquori M, Panno V. Hypertension and other risk factors in peripheral arterial disease. *Clin Exp Hypertens*. 1993;15 Suppl 1:71-89.
- 46 Islam TM, Fox CS, Mann D, Muntner P. Age-related associations of hypertension and diabetes mellitus with chronic kidney disease. *BMC Nephrol*. 2009 Jun 30;10:17.
- 47 Furberg AS, Thune I. Metabolic abnormalities (hypertension, hyperglycemia and overweight), lifestyle (high energy intake and physical inactivity) and endometrial cancer risk in a Norwegian cohort. *Int J Cancer*. 2003 May 10;104(6):659-76.
- 48 Jennings JR, Muldoon MF, Ryan C, Price JC, Greer P, Sutton-Tyrrell K, van der Veem FM, Meltzer CC. Reduced cerebral blood flow response and compensation among patients with untreated hypertension. *Neurology*. 2005 Apr 26;64(8):1358-65.
- 49 Jennings JR, Muldoon MF, Price J, Christie IC, Meltzer CC. Cerebrovascular support for cognitive processing in hypertensive patients is altered by blood pressure treatment. *Hypertension*. 2008 Jul;52(1):65-71.
- 50 Kuller LH, Margolis KL, Gaussoin SA, Bryan RN, Kerwin D, Limacher M, Wassertheil-Smoller S, Williamson J, Robinson JG. Relationship of hypertension, blood pressure, and blood pressure control with white matter abnormalities in the Women's Health Initiative Memory Study (WHIMS)-MRI trial. *J Clin Hypertens* (Greenwich). 2010 Mar;12(3):203-12.
- 51 Waldstein SR, Manuck SB, Ryan CM, Muldoon MF. Neuropsychological correlates of hypertension: review and methodologic considerations. *Psychol Bull*. 1991 Nov;110(3):451-68.
- 52 Foroughan M, Farahani ZG, Shariatpanahi M, Vaezinejad M, Kameran AA, Sheikhvatan M. Risk factors of Alzheimer's disease among Iranian population. *Curr Alzheimer Res*. 2008 Feb;5(1):70-2.
- 53 Redon J, Cifkova R, Laurent S, Nilsson P, Narkiewicz K, Erdine S, Manca G. Mechanisms of hypertension in the cardiometabolic syndrome. *J Hypertens*. 2009 Mar;27(3):441-51.
- 54 Li S, He H, Ding M, He C. The correlation of osteoporosis to clinical features: a study of 4382 female cases of a hospital cohort with musculoskeletal symptoms in southwest China. *BMC Musculoskelet Disord*. 2010 Aug 16;11:183.
- 55 Hyman L, Schachat AP, He Q, Leske MC. Hypertension, cardiovascular disease, and age-related macular degeneration. Age-Related Macular Degeneration Risk Factors Study Group. *Arch Ophthalmol*. 2000 Mar;118(3):351-8.
- 56 (1)Chew KK, Bremner A, Jamrozik K, Earle C, Stuckey B. Male erectile dysfunction and cardiovascular disease: is there an intimate nexus? *J Sex Med*. 2008 Apr;5(4):928-34.
- 57 Shibahara N, Matsuda H, Umeno K, Shimada Y, Itoh T, Terasawa K. The responses of skin blood flow, mean arterial pressure and R-R interval induced by cold stimulation with cold wind and ice water. *J Auton Nerv Syst*. 1996 Nov 6;61(2):109-15.
- 58 Dickinson BD, Havas S. Reducing the population burden of cardiovascular disease by reducing sodium intake: a report of the Council on Science and Public Health. *Arch Intern Med*. 2007 Jul 23;167(14):1460-8.
- 59 Morgenstern LB, Escobar JD, Sanchez BN, Hughes R, Zuniga BG, Garcia N, Lisabeth LD. Fast food and neighborhood stroke risk. *Ann Neurol*. 2009 Aug;66(2):165-70.
- 60 <http://nutritiondata.self.com/>
- 61 Zuccarelli MT, Faraj L. Sodium and potassium content of some fresh, frozen and canned vegetables. *Arch Latinoam Nutr*. 1986 Sep;36(3):477-82.
- 62 Anderson CA, Appel LJ, Okuda N, Brown JI, Chan Q, Zhao L, Ueshima H, Kesteloot H, Miura K, Curb JD, Yoshita K, Elliott P, Yamamoto ME, Stamler J. Dietary sources of sodium in China, Japan, the United Kingdom, and the United States, women and men aged 40 to 59 years: the INTERMAP study. *J Am Diet Assoc*. 2010 May;110(5):736-45.
- 63 Song J, Hu X, Shi M, Knepper MA, Ecelbarger CA. Effects of dietary fat, NaCl, and fructose on renal sodium and water transporter abundances and systemic blood pressure. *Am J Physiol Renal Physiol*. 2004 Dec;287(6):F1204-12.
- 64 Preuss HG. Diet, genetics and hypertension. *J Am Coll Nutr*. 1997 Aug;16(4):296-305.
- 65 Harshfield GA, Dong Y, Kapuku GK, Zhu H, Hanevol CD. Stress-induced sodium retention and hypertension: a review and hypothesis. *Curr Hypertens Rep*. 2009 Feb;11(1):29-34.
- 66 Shi Z, Yuan B, Taylor AW, Dai Y, Pan X, Gill TK, Wittert GA. Monosodium glutamate is related to a higher increase in blood pressure over 5 years: findings from the Jiangsu Nutrition Study of Chinese adults. *J Hypertens*. 2011 May;29(5):846-53.
- 67 Han TS, Gates E, Truscott E, Lean ME. Clothing size as an indicator of adiposity, ischaemic heart disease and cardiovascular risks. *J Hum Nutr Diet*. 2005 Dec;18(6):423-30.
- 68 Chockalingam A. Healthy weight - healthy blood pressure. *Can J Cardiol*. 2010 May;26(5):259-60.
- 69 Bellentani S, Scaglioni F, Marino M, Bedogni G. Epidemiology of non-alcoholic fatty liver disease. *Dig Dis*. 2010;28(1):155-61.
- 70 Ohashi K, Ouchi N, Matsuzawa Y. Adiponectin and Hypertension. *Am J Hypertens*. 2011 Mar;24(3):263-9.
- 71 Houston DK, Driver KE, Bush AJ, Kritchevsky SB. The association between cheese consumption and cardiovascular risk factors among adults. *J Hum Nutr Diet*. 2008 Apr;21(2):129-40.
- 72 Koi-Ramirez E, Castillo-Martinez L, Orea-Tejada A, Villa Romero AR, Vergara Castaneda A, Asensio Lafuente E. Waist circumference and fat intake are associated with high blood pressure in Mexican children aged 8 to 10 years. *J Am Diet Assoc*. 2009 Jun;109(6):996-1003.
- 73 Yang G, Shu XO, Gao YT, Zhang X, Li H, Zheng W. Impacts of weight change on prehypertension in middle-aged and elderly women. *Int J Obes (Lond)*. 2007 Dec;31(12):1818-25.
- 74 McCarron DA, Reusser ME. Body weight and blood pressure regulation. *Am J Clin Nutr*. 1996 Mar;63(3 Suppl):423S-425S.
- 75 Menghetti E, D'Addesa D, Censi L, Spagnolo A, Martone D, Cellitri R, Sette S. Hypertension in schoolchildren: research carried out in a secondary school in Rome and observations on dietary patterns. *Minerva Pediatr*. 2004 Jun;56(3):311-6.
- 76 Agadzhanov SA. Diet therapy of patients with chronic renal failure in its initial stage. *Vopr Pitan*. 1984 Nov-Dec;(6):28-31.
- 77 Wang YF, Yancy WS Jr, Yu D, Champagne C, Appel LJ, Lin PH. The relationship between dietary protein intake and blood pressure: results from the PREMIER study. *J Hum Nutr Diet*. 2008 Nov;22(11):745-54.
- 78 Rivas M, Garay RP, Escanero JF, Cia P Jr, Cia P, Alda JO. Soy milk lowers blood pressure in men and women with mild to moderate essential hypertension. *J Nutr*. 2002 Jul;132(7):1900-2.
- 79 Palanisamy N, Viswanathan P, Ravichandran MK, Anuradha CG. Renoprotective and blood pressure-lowering effect of dietary soy protein via protein kinase C beta 1 inhibition in a rat model of metabolic syndrome. *Can J Physiol Pharmacol*. 2010 Jan;88(1):28-37.
- 80 Nevala R, Vaskonen T, Vehniäinen J, Korpela R, Vapaatalo H. Soy based diet attenuates the development of hypertension when compared to casein based diet in spontaneously hypertensive rat. *Life Sci*. 2000;66(2):115-24.
- 81 Farag NH, Vincent AS, McKee BS, Al'Absi M, Whitsett TL, Lovallo WR. Sex differences in the hemodynamic responses to mental stress: effect of caffeine consumption. *Psychophysiology*. 2006 Jul;43(4):337-43.
- 82 Riksen NP, Rongen GA, Smits P. Acute and long-term cardiovascular effects of coffee: implications for coronary heart disease. *Pharmacol Ther*. 2009 Feb;121(2):185-91.
- 83 Jee SH, He J, Whelton PK, Suh I, Klag MJ. The effect of chronic coffee drinking on blood pressure: a meta-analysis of controlled clinical trials. *Hypertension*. 1999 Feb;33(2):647-52.
- 84 Langer S, Marshall LJ, Day AJ, Morgan MR. Flavonols and methylxanthines in commercially available dark chocolate: a study of the correlation with nonfat cocoa solids. *J Agric Food Chem*. 2011 Aug 10;59(15):8435-41.
- 85 Grassi D, Necozione S, Lippi C, Croce G, Valeri L, Pasqualetti P, Desideri G, Blumberg JB, Ferri C. Cocoa reduces blood pressure and insulin resistance and improves endothelium-dependent vasodilation in hypertensives. *Hypertension*. 2005 Aug;46(2):398-405.
- 86 Alonso A, de la Fuente C, Beunza JJ, Sánchez-Villegas A, Martínez-González MA. Chocolate consumption and incidence of hypertension. *Hypertension*. 2005 Dec;46(6):e21-2; author reply e22.
- 87 Dochi M, Sakata K, Oishi M, Tanaka K, Kobayashi E, Suwazono Y. Smoking as an independent risk factor for hypertension in a 14-year longitudinal study in male Japanese workers. *Tohoku J Exp Med*. 2009 Jan;217(1):37-43.
- 88 McGwin G, Liener J, Kennedy JI. Formaldehyde exposure and asthma in children: a systematic review. *Environ Health Perspect*. 2010 Mar;118(3):313-7.
- 89 Green MA, Egle Jr JI. Effects of intravenous acetaldehyde, acrolein, formaldehyde and propionaldehyde on arterial blood pressure following acute guanethidine treatment. *Res Commun Chem Pathol Pharmacol*. 1983 May;40(2):337-40.
- 90 Braune S, Wroclage C, Raczek J, Gailus T, Luking CH. Resting blood pressure increase during exposure to a radio-frequency electromagnetic field. *Lancet*. 1998 Jun 20;351(9119):1857-8.
- 91 Hiramatsu K, Yamada T, Katakura M. Acute effects of cold on blood pressure, renin-angiotensin-aldosterone system, catecholamines and adrenal steroids in man. *Clin Exp Pharmacol Physiol*. 1984 Mar-Apr;11(2):171-9.
- 92 Collins KJ. Low indoor temperatures and morbidity in the elderly. *Age Ageing*. 1986 Jul;15(4):212-20.
- 93 Shahar DR, Froom P, Harari G, Yerushalmi N, Lubin F, Kristal-Boneh E. Changes in dietary intake account for seasonal changes in cardiovascular disease risk factors. *Eur J Clin Nutr*. 1999 May;53(5):395-400.
- 94 Öpländer C, Volkmar CM, Paunel-Görgülü A, van Faassen EE, Heiss C, Kelm M, Halmer D, Mürtz M, Pallua N, Suschek CV. Whole body UVA irradiation lowers systemic blood pressure by release of nitric oxide from intracutaneous photolabile nitric oxide derivatives. *Circ Res*. 2009 Nov 6;105(10):1031-40.
- 95 Shani J, Kushlevsky AP, Harari M, Even-Paz Z. Sustained decrease of blood pressure in psoriatic patients during treatment at the Dead Sea. *Pharmacol Res*. 1995 Jun;31(6):355-9.
- 96 Kim MK, Il Kang M, Won Oh K, Kwon HS, Lee JH, Lee WC, Yoon KH, Son HY. The association of serum vitamin D level with presence of metabolic syndrome and hypertension in middle-aged Korean subjects. *Clin Endocrinol (Oxf)*. 2010 Sep;73(3):330-8.
- 97 Arakawa K. Hypertension and exercise. *Clin Exp Hypertens*. 1993 Nov;15(6):1171-9.
- 98 Demiot C, Dignat-George F, Fortrat JP, Sabatier F, Gharib C, Larina I, Gauquelin-Koch G, Hugbun R, Custaud MA. WISE 2005: chronic bed rest impairs microcirculatory endothelium in women. *Am J Physiol Heart Circ Physiol*. 2007 Nov;293(5):H3159-64.
- 99 Just H. Peripheral arterial disease and congestive heart failure: a review. *Am J Med*. 1991 May 29;90(5B):235-265.
- 100 Novo S, Pinto A, Alaimo G, Galati D, Strano A. Caloric flow and vascular resistance in borderline hypertensives in comparison with control subjects. *J Cardiovasc Pharmacol*. 1986;8 Suppl 5:S122-4.
- 101 Leung FP, Yung LM, Lahr I, Yao X, Chen ZY, Huang Y. Exercise, vascular wall and cardiovascular diseases: an update (Part 2). *Sports Med*. 2008;38(12):1009-24.
- 102 Yung LM, Lahr I, Yao X, Chen ZY, Huang Y, Leung FP. Exercise, vascular wall and cardiovascular diseases: an update (part 2). *Sports Med*. 2009;39(1):45-63.
- 103 Iwane M, Arita M, Tomimoto S, Satani O, Matsumoto M, Miyashita K, Nishio I. Walking 10,000 steps/day or more reduces blood pressure and sympathetic nerve activity in mild essential hypertension. *Hypertens Res*. 2000 Nov;23(6):573-80.
- 104 Mota MR, Padrono E, Lima LC, Arsa G, Bottaro M, Campbell CS, Simões HG. Effects of treadmill running and resistance exercises on lowering blood pressure during the daily work of hypertensive subjects. *J Strength Cond Res*. 2009 Nov;23(8):2331-8.
- 105 Meek SS. Effects of slow stroke back massage on relaxation in hospice clients. *Image J Nurs Sch*. 1993 Spring;25(1):17-21.
- 106 Nakawamoto R, Okamoto K, Yamada A, Oguni T. Effect of warm bathing on blood pressure in bedridden patients. *ippon Ronen Igakai Zasshi*. 1998 Apr;35(4):299-302.
- 107 Rafacz W, McGill SM. Wearing an abdominal belt increases diastolic blood pressure. *J Occup Environ Med*. 1996 Sep;38(9):925-7.
- 108 Wang SZ, Li S, Xu XY, Lin GP, Shao L, Zhao Y, Wang TH. Effect of slow abdominal breathing combined with biofeedback on blood pressure and heart rate variability in prehypertension. *J Altern Complement Med*. 2010 Oct;16(10):1039-45.
- 109 Jefferson LL. Exploring effects of therapeutic massage and patient teaching in the practice of diaphragmatic breathing on blood pressure, stress, and anxiety in hypertensive African-American women: an intervention study. *J Natl Black Nurses Assoc*. 2010 Jul;21(1):17-24.
- 110 Linde T, Sandhagen B, Iin C, Wikstr B, Danielson BG. Blood viscosity and peripheral vascular resistance in patients with untreated essential hypertension. *J Hypertens*. 1993 Jul;11(7):371-6.
- 111 Cicco G, Vicenti P, Stingi GD, Tarallo, Pirrelli A. Hemorheology in complicated hypertension. *Clin Hemorheol Microcirc*. 1999;21(3-4):315-9.
- 112 Bogar L. Hemorheology and hypertension: not "chicken or egg" but two chickens from similar eggs. *Clin Hemorheol Microcirc*. 2002;26(2):81-3.
- 113 Burstyn PG, Firth WR. Effects of three fat-enriched diets on the arterial pressure of rabbits. *Cardiovasc Res*. 1975 Nov;9(6):807-10.
- 114 Beegom R, Singh RB. Association of higher saturated fat intake with higher risk of hypertension in an urban population of Trivandrum in south India. *Int J Cardiol*. 1997 Jan 3;58(1):63-70.
- 115 Borghi C, Veronesi M, Cosentino E, Cicero AF, Kuria F, Dormi A, Ambrosioni E. Interaction between serum cholesterol levels and the renin-angiotensin system on the new onset of arterial hypertension in subjects with high-normal blood pressure. *J Hypertens*. 2007 Oct;25(10):2051-7.
- 116 Leviticus 7:23. King James Version of The Holy Bible.
- 117 Tamaya N, Uemura K, Yoshitaka S, Ueda M, Hattori A, Kuzuya M, Ohmoto Y, Muraguchi M, Nakamura J, Iguchi A. Noninvolvement of hypertriglyceridemia and hyperleptinemia in blood pressure increases induced by dietary lard in rats. *Drugs Exp Clin Res*. 2001;27(5-6):177-84.
- 118 Tamaya-Mori N, Uemura K, Tanaka S, Iguchi A. Aging accelerates dietary lard-induced increase in blood pressure in rats. *Exp Gerontol*. 2003 Aug;38(8):905-10.
- 119 Deuteronomy 14:8. King James Version of The Holy Bible.
- 120 Slavicek J, Kittnar O, Dohnalová A, Trojan S, Novák V, Tichý JA, Trefný ZM. Effect of a 10-day animal fat-free diet on cholesterol and glucose serum levels, blood pressure and body weight in 50-year-old volunteers. *Sb Lek*. 2001;102(4):519-25.
- 121 Stolberg HO, Norman G, Trop I. Randomized controlled trials. *AJR Am J Roentgenol*. 2004 Dec;183(6):1539-44.
- 122 Daniel 1:12. King James Version of The Holy Bible.
- 123 Naito Y, Yoshida H, Nagata T, Tanaka A, Ono H, Ohara N. Dietary intake of rapeseed oil or soybean oil as the only fat nutrient in spontaneously hypertensive rats and Wistar Kyoto rats - blood pressure and pathophysiology. *Toxicology*. 2000 May 5;146(2-3):197-208.
- 124 Kaufman LN, Peterson MM, Smith SM. Hypertensive effect of polyunsaturated dietary fat. *Metabolism*. 1994 Jan;43(1):1-3.
- 125 Naito Y, Nagata T, Takano Y, Nagatsu T, Ohara N. Rapeseed oil ingestion and exacerbation of hypertension-related conditions in stroke prone spontaneously hypertensive rats. *Toxicology*. 2003 May 3;187(2-3):205-16.
- 126 Sorriguer F, Rojo-Martinez G, Dobarganes MC, Garcia Almeida JM, Esteve I, Beltan M, Ruiz De Adana MS, Tinahones F, Gomez-Zumaguer JM, Garcia Fuentes E, Gonzalez-Romero S. Hypertension is related to the degradation of dietary frying oils. *Am J Clin Nutr*. 2003 Dec;78(6):1092-7.
- 127 Gosmanov AR, Smiley DD, Robalino G, Siquiera J, Khan B, Le NA, Patel RS, Quyuyumi AA, Peng L, Kitabchi AE, Umpierrez GE. Effects of oral and intravenous fat load on blood pressure, endothelial function, sympathetic activity, and oxidative stress in obese healthy subjects. *Am J Physiol Endocrinol Metab*. 2010 Dec;299(6):E953-8.
- 128 Giannotti G, Doerries C, Mochalra PS, Mueller MF, Bahlmann FH, Horvath T, Jiang H, Sorrentino SA, Steenken N, Manes C, Marzilli M, Rudolph KL, Luscher TF, Drexlher H, Landmesser U. Impaired endothelial repair capacity of early endothelial progenitor cells in prehypertension: relation to endothelial dysfunction. *Hypertension*. 2010 Jun;55(6):1389-97.
- 129 Cugini P, Baldoni F, De Rosa R, Pandolfi C, Colotto M, Buccarella PA, Zamparelli C, Berti D, Passini B, Roncoroni V, Sabino D, Capria A. Higher blood pressure load (baric impact) in normotensives with endothelial dysfunction: a pathophysiological status of "pre-hypertension". *Clin Ter*. 2002 Sep-Oct;153(5):309-15.
- 130 Bleande MC, Bard M, Sowers JR, Winer N. High-fat meal impairs vascular compliance in a subgroup of young healthy subjects. *Metabolism*. 2005 Oct;54(10):1337-44.
- 131 Sofola O, Knill A, Myers D, Hainsworth R, Drinkhill M. High-salt diet and responses of the pressurized mesenteric artery of the dog to noradrenaline and acetylcholine. *Clin Exp Pharmacol Physiol*. 2004 Oct;31(10):696-9.
- 132 Fiore MC, Jimenez PM, Cremonesi D, Juncos LI, Garcia NH. Statins reverse renal inflammation and endothelial dysfunction induced by chronic high salt intake. *Am J Physiol Renal Physiol*. 2011 Aug;301(2):F263-70.
- 133 Lind L. Lipids and endothelium-dependent vasodilation—a review. *Lipids*. 2002 Jan;37(1):1-15.
- 134 Moriel P, Sevastian A, Ajzen S, Zanella MT, Plavnik FL, Rubbo H, Abdalla DS. Nitric oxide, cholesterol oxides and endothelium-dependent vasodilation in plasma of patients with essential hypertension. *Braz J Med Biol Res*. 2002 Nov;35(11):1301-9.
- 135 Sasaki S, Higashi Y, Nakagawa K, Kimura M, Noma K, Sasaki S, Hara K, Matsuura H, Goto C, Oshima T, Chayama K. A low-calorie diet improves endothelium-dependent vasodilation in obese patients with essential hypertension. *Am J Hypertens*. 2002 Apr;15(4 Pt 1):302-9.
- 136 Vaag A, Bruns C, Appel JS, Touros S. Metabolic consequences of overeating. *Ugeskr Laeger*. 2006 Jan 9;168(2):183-7.
- 137 Gomes MB, Afonso FS, Cailleaux S, Almeida AL, Pinto LF, Tiberiçá E. Glucose levels observed in daily clinical practice induce endothelial dysfunction in the rabbit macro- and microcirculation. *Fundam Clin Pharmacol*. 2004 Jun;18(3):339-46.
- 138 Tran LT, Yuen VG, McNeill JH. The fructose-fed rat: a review on the mechanisms of fructose-induced insulin resistance and hypertension. *Mol Cell Biochem*. 2009 Dec;332(1-2):145-59.
- 139 Winer N, Sowers JR. Vascular compliance in diabetes. *Curr Diab Rep*. 2003 Jun;3(3):230-4.
- 140 Lin CL, Fang TC, Gueng MK. Vascular dilatory functions of ovo-lactovegetarians compared with omnivores. *Atherosclerosis*. 2001 Sep;158(1):247-51.
- 141 Katz DL, Nawaz H, Kulkarni J, Ganamovore V, Chan W, Ahmedi R, Sarrel PM. Acute effects of oats and vitamin E on endothelial responses to ingested fat. *Am J Prev Med*. 2001;20(2):114-9.
- 142 Suganuma H, Inakuma T. Protective effect of dietary tomato against endothelial dysfunction in hypercholesterolemic mice. *Biosci Biotechnol Biochem*. 1999 Jan;63(1):78-82.
- 143 Sato J, O'Brien T, Katsic ZS, Fu A, Nygren J, Singh R, Nair KS. Dietary antioxidants preserve endothelium dependent vasorelaxation in overfed rats. *Atherosclerosis*. 2002 Apr;161(2):327-33.
- 144 Ribeiro Jorge PA, Neyra LC, Ozaki RM, de Almeida E. Improvement in the endothelium-dependent relaxation in hypercholesterolemic rabbits treated with vitamin E. *Atherosclerosis*. 1998 Oct;140(2):333-9.
- 145 Browning JD, Reeves PG, O'Dell BL. Zinc deficiency in rats reduces the vasodilation response to bradykinin and prostacyclin. *J Nutr*. 1987 Mar;117(3):490-5.
- 146 Schuschke DA, Saari JT, Miller FN. A role for dietary copper in nitric oxide-mediated vasodilation. *Microcirculation*. 1995 Dec;2(4):371-6.
- 147 Covington MB. Omega-3 fatty acids. *Am Fam Physician*. 2004 Jul 1;70(1):133-40.
- 148 Begg DP, Sinclair AJ, Stahl LA, Premaratna SD, Hafandi A, Jois M, Weisinger RS. Hypertension induced by omega-3 polyunsaturated fatty acid deficiency is alleviated by alpha-linolenic acid regardless of dietary source. *Hypertens Res*. 2010 Aug;33(8):808-13.
- 149 Zhang HY, Reddy S, Kotchen TA. A high sucrose, high linoleic acid diet potentiates hypertension in the Dahl salt sensitive rat. *Am J Hypertens*. 1999 Feb;12(2 Pt 1):183-7.
- 150 Ahrens RA, Rejto D, Walker, Am J Clin Nutr. 1975 Mar;28(3): 197-200.
- 151 United States Department of Agriculture, Office of Communications, Agriculture Fact Book 2001-2002, ISBN 001-000-04709-4.
- 152 Peteri-Petridi I. High glucose and renin release: the role of succinate and GPR91. *Kidney Int*. 2010 Dec;78(12):1214-7.
- 153 Meyer KA, Kushi LH, Jacobs DR Jr, Slavin J, Sellers TA, Folsom AR. Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. *Am J Clin Nutr*. 2000 Apr;71(4):921-30.
- 154 Pavan L, Casiglia E, Braga LM, Winnick M, Puato M, Paultetto P, Pessina AC. Effects of a traditional lifestyle on the cardiovascular risk profile: the Amondava population of the Brazilian Amazon. Comparison with matched African, Italian and Polish populations. *J Hypertens*. 1999 Jun;17(6):749-56.
- 155 Preuss HG, Fournier RD, Chieuh CC, Kopin IJ, Knappa J, DiPette D, More NS, Rao NA. Refined carbohydrates affect blood pressure and retinal vasculature in spontaneously hypertensive and Wistar-Kyoto rats. *J Hypertens Suppl*. 1986 Oct;4(3):S459-62.
- 156 Chen L, Caballero B, Mitchell DC, Loria C, Lin PH, Champagne CM, Elmer PJ, Ard JD, Batich BC, Anderson CA, Appel LJ. Reducing consumption of sugar-sweetened beverages is associated with reduced blood pressure: a prospective study among United States adults. *Circulation*. 2010 Jun 8;121(22):2398-406.
- 157 Ziemian SJ, Kass DA. Advanced glycation endproduct crosslinking in the cardiovascular system: potential therapeutic target for cardiovascular disease. *Drugs*. 2004;64(5):459-70.
- 158 Vasdes V, Gill V, Singal P. Role of advanced glycation endproducts in hypertension and atherosclerosis: therapeutic implications. *Clin Biochem Biophys*. 2007;49(1):48-63.
- 159 Anderson RA. Chromium and insulin resistance. *Nutr Res Rev*. 2003 Dec;16(2):267-75.
- 160 Rupp H. Insulin resistance, hyperinsulinemia, and cardiovascular disease. The need for novel dietary prevention strategies. *Basic Res Cardiol*. 1992 Mar-Apr;87(2):99-105.

165 Torimitsu M, Nagase R, Yanagi M, Homma M, Sasai Y, Ito Y, Hayamizu K, Nonaka S, Hosono T, Kise M, Seki T, Ariga T. Replacing white rice with pre-germinated brown rice mildly ameliorates hyperglycemia and imbalance of adipocytokine levels in type 2 diabetes model rats. *J Nutr Sci Vitaminol (Tokyo)*. 2010;56(5):287-92.

166 Radhika G, Van Dam RM, Sudha V, Ganesan A, Mohan V. Refined grain consumption and the metabolic syndrome in urban Asian Indians (Chennai Urban Rural Epidemiology Study 57). *Metabolism*. 2009 May;58(5):675-81.

167 Kynde I, Johnsen NF, Wedderkopp N, Bygberg JE, Helge JW, Heitmann BL. Intake of total dietary sugar and fibre is associated with insulin resistance among Danish 8-10- and 14-16-year-old girls but not boys. *European Youth Heart Studies I and II. Public Health Nutr*. 2010 Oct;13(10):1669-74.

168 Ferreira Medel R, Lombardo VB, Chicho A. β -Cell adaptation/dysfunction in an animal model of dyslipidemia and insulin resistance induced by the chronic administration of a sucrose-rich diet. *Islets*. 2010 Nov 12;2(6):367-73.

169 Dekker MJ, Su Q, Baker C, Rutledge AC, Adeli K. Fructose: a highly lipogenic nutrient implicated in insulin resistance, hepatic steatosis, and the metabolic syndrome. *Am J Physiol Endocrinol Metab*. 2010 Nov;299(5):E685-94.

170 Stanhope KL, Schwarz JM, Keim NL, Griffen SC, Bremer AA, Graham JL, Hatcher B, Cox CL, Dyachenko A, Zhang W, McGahan JP, Seibert A, Krauss RM, Chiu S, Schaefer EJ, Ai M, Otokozawa S, Nakajima K, Nakano T, Beyens C, Hellerstein MK, Berglund L, Havel PJ. Consuming fructose-sweetened, not glucose-sweetened, beverages increases visceral adiposity and lipids and decreases insulin sensitivity in overweight/obese humans. *J Clin Invest*. 2009 May;119(5):1322-34.

171 Costa RR, Villela NR, Souza MD, Boa BC, Cyrino FZ, Silva SV, Lisboa PC, Moura EG, Barja-Fidalgo TC, Bouckea E. High fat diet induces central obesity, insulin resistance and microvascular dysfunction in hamsters. *Microvasc Res*. 2011 Aug 25.

172 Bruns W. Treatment of type 2 (non-insulin dependent) diabetes and the metabolic syndrome with diet. *Z Gesamte Inn Med*. 1991 Oct;46(15):563-7.

173 Reaven GM. Insulin resistance: the link between obesity and cardiovascular disease. *Med Clin North Am*. 2011 Sep;95(5):875-92.

174 Robins SJ, Lyass A, Zachariah JP, Massaro JM, Vasan RS. Insulin resistance and the relationship of a dyslipidemia to coronary heart disease: the Framingham Heart Study. *Arterioscler Thromb Vasc Biol*. 2011 May;31(5):1208-14.

175 Lai TS, Curhan GC, Forman JP. Insulin resistance and risk of incident hypertension among men. *J Clin Hypertens (Greenwich)*. 2009 Sep;11(9):483-90.

176 Sung KC, Lim S, Rosenson RS. Hyperinsulinemia and homeostasis model assessment of insulin resistance as predictors of hypertension: a 5-year follow-up study of Korean sample. *Am J Hypertens*. 2011 Sep;24(9):1041-5.

177 Kawamoto R, Kohara K, Tabara Y, Abe M, Kusunoki T, Miki T. Insulin resistance and prevalence of prehypertension and hypertension among community-dwelling persons. *J Atheroscler Thromb*. 2010 Feb 26;17(2):148-55.

178 Harter AL, McTernan PG, McTernan CL, Crocker J, Starczynski J, Barnett AH, Matyka K, Kumar S. Insulin increases angiotensinogen expression in human abdominal subcutaneous adipocytes. *Diabetes Obes Metab*. 2003 Nov;5(6):462-7.

179 Jalal DI, Smits G, Johnson RJ, Chonchol M. Increased Fructose Associates with Elevated Blood Pressure. *J Am Soc Nephrol*. 2010 Sep;21(9):1543-9.

180 Winkelmayer WC, Stampfer MJ, Willett WC, Curhan G. Habitual caffeine intake and the risk of hypertension in women. *JAMA*. 2005 Nov 9;294(18):2330-5.

181 Roberts JH. Overlooked aspartame-induced hypertension. *South Med J*. 2008 Sep;101(9):969.

182 Ulbricht C, Isaac R, Milkin T, Poole EA, Rusie E, Grimes Serrano JM, Weissner W, Windsor RC, Woods J. An evidence-based systematic review of stevia by the Natural Standard Research Collaboration. *Cardiovasc Hematol Agents Med Chem*. 2010 Apr;8(2):1-13.

183 Lee CN, Wong KL, Liu JC, Chen YJ, Cheng JT, Chan P. Inhibitory effect of stevioside on calcium influx to produce antihypertension. *Planta Med*. 2001 Dec;67(9):796-9.

184 Peralta CA, Adeney KL, Shlipak MG, Jacobs DR Jr, Duprez D, Blumenthal D, Polak J, Psaty B, Kestenbaum BR. Structural and functional vascular alterations and incident hypertension in normotensive adults: the Multi-Ethnic Study of Atherosclerosis. *Am J Epidemiol*. 2010 Jan 1;171(1):63-71.

185 Al-Nimer MS, Hussein II, Lasso WS. Subtype of hypertension is evidence for preclinical atherosclerosis. A study of carotid artery ultrasonography and biochemical markers. *Neurosciences (Riyadh)*. 2010 Apr;15(2):79-83.

186 Floras JS. Hypertension, sleep apnea, and atherosclerosis. *Hypertension*. 2009 Jan;53(1):1-3.

187 Li JJ, Chen JL. Inflammation may be a bridge connecting hypertension and atherosclerosis. *Med Hypotheses*. 2005;64(5):925-9.

188 Mattace-Raso FU, Verwoert GC, Hofman A, Witteman JC. Inflammation and incident-isolated systolic hypertension in older adults: the Rotterdam study. *J Hypertens*. 2010 May;28(5):892-5.

189 Zhang Y, Thompson AM, Tong W, Xu T, Chen J, Zhao L, Kelly TN, Chen CS, He J. Biomarkers of inflammation and endothelial dysfunction and risk of hypertension among Inner Mongolians in China. *J Hypertens*. 2010 Jan;28(1):35-40.

190 Bussemaker E, Hillebrand U, Hausberg M, Pavenstadt H, Oberleitner H. Pathogenesis of hypertension: interactions among sodium, potassium, and aldosterone. *Am J Kidney Dis*. 2010 Jun;55(6):1111-20.

191 MacGregor GA, Smith SD, Markandya ND, Sagnella GA. Does increasing potassium intake lower blood pressure in essential hypertension? *J Cardiovasc Pharmacol*. 1984;6:1524-9.

192 Haddy FJ. Role of dietary salt in hypertension. *Life Sci*. 2006 Sep 20;79(17):1585-92.

193 Lenda DM, Boegehold MA. Effect of a high salt diet on microvascular antioxidant enzymes. *J Vasc Res*. 2002 Jan-Feb;39(1):41-50.

194 Chandramohan G, Bai Y, Norris K, Rodriguez-Turbe B, Vaziri ND. Effects of dietary salt on intrarenal angiotensin system, NAD(P)H oxidase, COX-2, MCP-1 and PAI-1 expressions and NF-kappaB activity in salt-sensitive and -resistant rat kidneys. *Am J Nephrol*. 2008;28(1):158-67.

195 Navas-Acien A, Guallar E, Silbergeld EK, Rothenberg SJ. Lead exposure and cardiovascular disease—a systematic review. *Environ Health Perspect*. 2007 Mar;115(3):472-82.

196 Kwok RK, Mendola P, Liu ZY, Sawitz DA, Heiss G, Ling HL, Xia Y, Lobdell D, Zeng D, Thorp JM Jr, Creason JP, Mumford JL. Drinking water arsenic exposure and blood pressure in healthy women of reproductive age in Inner Mongolia, China. *Toxicol Appl Pharmacol*. 2007 Aug 1;222(3):337-43.

197 Ross EA, Szabo NJ, Tebbett IR. Lead content of calcium supplements. *JAMA*. 2000 Sep 20;284(11):1425-9.

198 Chen CJ, Wang SL, Chiou JM, Tseng CH, Chiou HY, Hsueh YM, Chen SY, Wu MM, Lai MS. Arsenic and diabetes and hypertension in human populations: a review. *Toxicol Appl Pharmacol*. 2007 Aug 1;222(3):298-304.

199 Sullivan MJ, Leavey S. Heavy metals in bottled natural spring water. *J Environ Health*. 2011 Jun;73(10):18-33.

200 Taylor DA. Funky chicken: consumers exposed to arsenic in poultry. *Environ Health Perspect*. 2004 Jan;112(1):A50.

201 Holman A, Sibbily V. Arsenic residues in eggs from laying hens fed with a diet containing arsenic (III) oxide. *Arch Environ Contam Toxicol*. 1997 May;32(4):407-10.

202 Lasky T, Sun W, Kadry A, Hoffman MK. Mean total arsenic concentrations in chicken 1989-2000 and estimated exposures for consumers of chicken. *Environ Health Perspect*. 2004 Jan;112(1):18-21.

203 Choi BS, Choi SJ, Kim DW, Huang M, Kim NY, Park KS, Kim CY, Lee HM, Yum UN, Han ES, Kang TS, Yu JJ, Park JD. Effects of repeated seafood consumption on urinary excretion of arsenic species by volunteers. *Arch Environ Contam Toxicol*. 2010 Jan;58(1):222-9.

204 Wu L, Noyan Ashraf MH, Facci M, Wang R, Paterson PG, Ferrie A, Juurlink BH. Dietary approach to attenuate oxidative stress, hypertension, and inflammation in the cardiovascular system. *Proc Natl Acad Sci U S A*. 2004 May 4;101(18):7059-9.

205 Seaman DR. The diet-induced proinflammatory state: a cause of chronic pain and other degenerative diseases? *J Manipulative Physiol Ther*. 2002 Mar-Apr;25(3):168-79.

206 Azadbakht L, Surkan PJ, Esmailzadeh A, Willett WC. The Dietary Approaches to Stop Hypertension eating plan affects C-reactive protein, coagulation abnormalities, and hepatic function tests among type 2 diabetic patients. *J Nutr*. 2011 Jun;141(6):1083-8.

207 Ceriello A, Giugliano D, Quattraro A, Lefebvre PJ. Anti-oxidants show an anti-hypertensive effect in diabetic and hypertensive subjects. *Clin Sci (Lond)*. 1991 Dec;81(6):739-42.

208 Agarwal D, Haque M, Sriaramula S, Mariappan N, Pariat R, Francis J. Role of proinflammatory cytokines and redox homeostasis in exercise-induced delayed progression of hypertension in spontaneously hypertensive rats. *Hypertension*. 2009 Dec;54(6):935-40.

209 Desvarieux M, DeMeer RT, Jacobs DR Jr, Rudek T, Boden-Albala B, Sacco RL, Papaianou PN. Periodontal bacteria and hypertension: the oral infections and vascular disease epidemiology study (INVEST). *J Hypertens*. 2010 Jul;28(7):1413-21.

210 Paulis L, Pechanova O, Zicha J, Barta A, Gardlik R, Celec P, Kunes J, Simko F. Melatonin interactions with blood pressure and vascular function during L-NAME-induced hypertension. *J Pineal Res*. 2010 Mar;48(2):102-8.

211 Reiter RJ, Korkmaz A. Clinical aspects of melatonin. *Saudi Med J*. 2008 Nov;29(11):1537-47.

212 Gooley JF, Chamberlain K, Smith KA, Khalsa SB, Rajaratnam SM, Van Reen E, Zeitzer JM, Czeisler CA, Lockley SW. Exposure to Room Light before Bedtime Suppresses Melatonin Onset and Shortens Melatonin Duration in Humans. *J Clin Endocrinol Metab*. 2011 Mar;96(3):E463-72.

213 Simko F, Pechanova O, Pelouch V, Krabiciovicova K, Celec P, Palffy R, Bednarova K, Vrankova S, Adamcova M, Paulis L. Continuous light and L-NAME-induced left ventricular remodeling: different protection with melatonin and captopril. *J Hypertens*. 2010 Sep;28 Suppl 1:S13-8.

214 Forman JP, Curhan GC, Schernhammer ES. Urinary melatonin and risk of incident hypertension among young women. *J Hypertens*. 2010 Mar;28(3):446-51.

215 Reiter RJ, Tan DX, Korkmaz A. The circadian melatonin rhythm and its modulation: possible impact on hypertension. *J Hypertens Suppl*. 2009 Aug;27(6):S17-20.

216 McCubbin JA, Pilcher JJ, Moore DD. Blood Pressure Increases During a Simulated Night Shift in Persons at Risk for Hypertension. *Int J Behav Med*. 2010 Dec;17(4):314-20.

217 Su TC, Lin LY, Baker D, Schall PL, Chen MF, Huang WC, Chen CF, Wang JD. Elevated blood pressure, decreased heart rate variability and incomplete blood pressure recovery after a 12-hour night shift work. *J Occup Health*. 2008;50(5):380-6.

218 Calhoun DA, Harding SM. Sleep and hypertension. *Chest*. 2010 Aug;138(2):434-43.

219 Steffen ML, Kroenke CH, Yu X, Pereira MA, Slattery ML, Van Horn L, Gross MD, Jacobs DR Jr. Associations of plant food, dairy product, and meat intakes with 15-year incidence of elevated blood pressure in young black and white adults: the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Am J Clin Nutr*. 2005 Dec;82(6):1169-77.

220 Burch GE. Pork and hypertension. *Am Heart J*. 1973 Nov;86(5):713-4.

221 Leviticus 11:7 King James Version of The Holy Bible.

222 Ivan A, Gröll M, Duda R, Hurlui J, Serban F, Mitroi I. Interventions associated with some risk factors for essential arterial hypertension. I. Epidemiologic observations in adults. *Rev Med Chir Soc Med Nat Iasi*. 1989 Apr-Jun;93(2):309-13.

223 Polovitchina OV, Oshchepkova EV, Dmitriev VA, Titov VN. Role of uric acid in development of essential hypertension: modern conceptions. *Ter Arkh*. 2011;83(8):38-41.

224 Zhang L, Curhan GC, Forman JP. Diet-dependent net acid load and risk of incident hypertension in United States women. *Hypertension*. 2009 Oct;54(4):751-5.

225 Remer T, Manz F. Potential renal acid load of foods and its influence on urine pH. *J Am Diet Assoc*. 1995 Jul;95(7):791-7.

226 Mellen PB, Bleyer AJ, Erlinger TP, Evans GW, Nieto FJ, Wagenknecht LE, Wofford MR, Herrington DM. Serum uric acid predicts incident hypertension in a biethnic cohort: the atherosclerosis risk in communities study. *Hypertension*. 2006 Dec;48(6):1037-42.

227 Chuang SY, Lee SC, Hsieh YT, Pan WH. Trends in hyperuricemia and gout prevalence: Nutrition and Health Survey in Taiwan from 1993-1996 to 2005-2008. *Asia Pac J Clin Nutr*. 2011;20(2):301-8.

228 Villegas R, Xiang YB, Elasy T, Xu WH, Cai H, Cai Q, Linton MF, Fazio S, Zheng W, Shu XO. Purine-rich foods, protein intake, and the prevalence of hyperuricemia: The Shanghai Men's Health Study. *Nutr Metab Cardiovasc Dis*. 2011 Jan 27.

229 Nguyen S, Choi HK, Lustig RH, Hsu CY. Sugar-sweetened beverages, serum uric acid, and blood pressure in adolescents. *J Pediatr*. 2009 Jun;154(6):807-13.

230 Ka T, Moriwaki Y, Inokuchi T, Yamamoto A, Takahashi S, Tsutsumi Z, Yamamoto T. Effects of allopurinol on beer-induced increases in plasma concentrations and urinary excretion of purine bases (uric acid, hypoxanthine, and xanthine). *Horm Metab Res*. 2006 Mar;38(3):188-92.

231 Aras B, Kafkazez N, Tuğcu V, Kemahli E, Ozyab B, Polat H, Taşci AN. Can lemon juice be an alternative to potassium citrate in the treatment of urinary calcium stones in patients with hypocitraturia? A prospective randomized study. *Urol Res*. 2008 Dec;36(6):313-7.

232 Jacob RA, Spinuzzi GM, Simon VA, Kelley DS, Prior RL, Hess-Pierce B, Kader AA. Consumption of cherries lowers plasma urate in healthy women. *J Nutr*. 2003 Jun;133(6):1826-9.

233 Tulipani S, Mezzetti B, Battino M. Impact of strawberries on human health: insight into originally discussed bioactive compounds for the Mediterranean diet. *Public Health Nutr*. 2009 Sep;12(9A):1656-62.

234 Pamplona-Roger G. ENCYCLOPEDIA OF FOODS AND THEIR HEALING POWER. Hagerstown Maryland, Review & Herald Publishing association, 2004.

235 Taylor B, Irving HM, Balunias D, Roerecke M, Patra J, Mohapatra S, Rehm J. Alcohol and hypertension: gender differences in dose-response relationships determined through systematic review and meta-analysis. *Addiction*. 2009 Dec;104(12):1981-90.

236 Ueshima H, Shimamoto T, Iida M, Konishi M, Tanigaki M, Doi M, Tsuboioka K, Nagano E, Tsuda C, Ozawa H, et al. Alcohol intake and hypertension among urban and rural Japanese populations. *J Chronic Dis*. 1984;37(7):585-92.

237 Proverbs 20:1, King James Version of the Holy Bible.

238 Teixeira Rde C, Molina Mdel C, Zandonade E, Mill JG. Cardiovascular risk in vegetarians and omnivores: a comparative study. *Arq Bras Cardiol*. 2007 Oct;89(4):237-44.

239 Carvalho J, Baruzzi RG, Howard PF, Poulter N, Alpers MP, Franco LJ, Marcopito LF, Spooner VJ, Dyer AR, Elliott P, Stamler J, Stamler R. Blood pressure in four remote populations in the INTERSALT Study. *Hypertension*. 1989 Sep;14(3):238-46.

240 Pavan L, Casiglia E, Braga LM, Winnick M, Puato M, Pualetto P, Pessina AC. Effects of a traditional lifestyle on the cardiovascular risk profile: the Amondava population of the Brazilian Amazon. Comparison with matched African, Italian and Polish populations. *J Hypertens*. 1999 Jun;17(6):749-56.

241 Appleby PN, Davey GK, Key TJ. Hypertension and blood pressure among meat eaters, fish eaters, vegetarians and vegans in EPIC-Oxford. *Public Health Nutr*. 2002 Oct;5(5):645-54.

242 Fraser GE. Vegetarian diets: what do we know of their effects on common chronic diseases? *Am J Clin Nutr*. 2009 May;89(5):1607S-1612S.

243 Lindahl O, Lindvall L, Spångberg A, Stenram A, Ockerman PA. A vegan regimen with reduced medication in the treatment of hypertension. *Br J Nutr*. 1984 Jul;52(1):11-20.

244 Alonso A, de la Fuente C, Martí-Arnuau AM, de Irala J, Martínez JA, Martínez-González MA. Fruit and vegetable consumption is inversely associated with blood pressure in a Mediterranean population with a high vegetable/fat intake: the Segumiento Universidad de Navarra (SUN) Study. *Br J Nutr*. 2004 Aug;92(2):311-9.

245 Dauchet L, Czernichow S, Bertrai S, Blacher J, Galan P, Hercberg S, SFHTA. Fruits and vegetables intake in the SU.VI.MAX study and systolic blood pressure change. *Arch Mal Coeur Vaiss*. 2006 Jul-Aug;99(7-8):669-73.

246 He K, Song Y, Belin RJ, Chen Y. Magnesium intake and the metabolic syndrome: epidemiologic evidence to date. *J Cardiometab Syndr*. 2006 Fall;1(5):351-5.

247 D. Tsi and B. K. H. Tan. Cardiovascular Pharmacology of 3-n-butylphthalide (found in celery) in Spontaneously Hypertensive Rats. *Phytotherapy Research*. Dec 1997; 11(8): 576-582.

248 Gilani AH, Shaheen E, Saaved SA, Bibi S, Irfanullah, Sadiq M, Faizi S. Hypotensive action of coumarin glycosides from *Daucus carota*. *Phytomedicine*. 2000 Oct;7(5):423-6.

249 Tapsell LC, Hemphill I, Cobiac L, Patch CS, Sullivan DR, Fenech M, Roodenrys S, Keogh JB, Clifton PM, Williams PG, Fazio VA, Inge KE. Health benefits of herbs and spices: the past, the present, the future. *Med J Aust*. 2006 Aug 21;185(4 Suppl):S4-24.

250 Wu L, Noyan Ashraf MH, Facci M, Wang R, Paterson PG, Ferrie A, Juurlink BH. Dietary approach to attenuate oxidative stress, hypertension, and inflammation in the cardiovascular system. *Proc Natl Acad Sci U S A*. 2004 May 4;101(18):7059-9.

251 Ferreira LF, Behnke BJ. A toast to health and performance! Beetroot juice lowers blood pressure and the O2 cost of exercise. *J Appl Physiol*. 2011 Mar;110(3):585-6.

252 Umar A, Imam G, Yimin W, Kerim P, Tohti I, Berke B, Moore N. Antihypertensive effects of *Cicum basilicum* L. (OBL) on blood pressure in renovascular hypertensive rats. *Hypertens Res*. 2010 Jul;33(7):727-30.

253 Sobenin IA, Andrianova IV, Fomchenkov IV, Gorchakova TV, Orekhov AN. Time-released garlic powder tablets lower systolic and diastolic blood pressure in men with mild and moderate arterial hypertension. *Hypertens Res*. 2009 Jun;32(6):433-7.

254 Kawamoto E, Sakai Y, Okamura Y, Yamamoto Y. Effects of boiling on the antihypertensive and antioxidant activities of onion. *J Nutr Sci Vitaminol (Tokyo)*. 2004 Jun;50(3):171-6.

255 Utsugi MT, Ohsue K, Tanaka T, Kikuya K, Kurimoto A, Sato R, Suzuki K, Metoki H, Arha A, Tsubono Y, Imai Y. Fruit and vegetable consumption and the risk of hypertension determined by self measurement of blood pressure at home: the Ohasama study. *Hypertens Res*. 2008 Jul;31(7):1435-43.

256 Kwon YJ, Apostolidis E, Shetty K. In vitro studies of eggplant (*Solanum melongena*) phenolics as inhibitors of key enzymes relevant for type 2 diabetes and hypertension. *Bioresour Technol*. 2008 May;99(8):2981-8.

257 Paran E, Novack V, Engelhard YN, Hazan-Halevy I. The effects of natural antioxidants from tomato extract in treated but uncontrolled hypertensive patients. *Cardiovasc Drugs Ther*. 2009 Apr;23(2):145-51.

258 Kwon YJ, Apostolidis E, Kim YC, Shetty K. Health benefits of traditional corn, beans, and pumpkin: in vitro studies for hyperglycemia and hypertension management. *J Med Food*. 2007 Jun;10(2):266-75.

259 Apostolidis E, Kwon YJ, Shetty K. Potential of cranberry-based herbal synergies for diabetes and hypertension management. *Asia Pac J Clin Nutr*. 2006;15(3):433-41.

260 Sasaki N. Life style and blood pressure: the protective effect of apple-eating habits on high blood pressure in a high-salt population. *Nippon Eisegaku Zasshi*. 1990 Dec;45(5):953-63.

261 Aviram M, Rosenbalt M, Gaitini D, Nitecki S, Hoffman A, Dornfeld L, Volkova N, Presser D, Attias J, Liker H, Hayek T. Pomegranate juice consumption for 3 years by patients with carotid artery stenosis reduces common carotid intima-media thickness, blood pressure and LDL oxidation. *Clin Nutr*. 2004 Jun;23(3):423-33.

262 Diaz-Juarez JA, Tenorio-López FA, Zarco-Olvera G, Valle-Monrealgrón LD, Torres-Narváez JC, Pastelín-Hernández G. Effect of *Citrus paradisi* extract and juice on arterial pressure both in vitro and in vivo. *Phytother Res*. 2009 Jul;23(7):948-54.

263 Gilani AH, Khan AU, Shah AJ, Connor J, Jabeen Q. Blood pressure lowering effect of olive is mediated through calcium channel blockade. *Int J Food Sci Nutr*. 2005 Dec;56(8):613-20.

264 Galan P, Vergnaud AC, Troulakis I, Buyck JF, Blacher J, Czernichow S, Hercberg S. Low total and nonheme iron intakes are associated with a greater risk of hypertension. *J Nutr*. 2010 Jan;140(1):75-80.

265 Krishna GG. Role of potassium in the pathogenesis of hypertension. *Am J Med Sci*. 1994 Feb;307 Suppl 1:S21-5.

266 Dyckner T, Wester PO. Potassium/magnesium depletion in patients with cardiovascular disease. *Am J Med*. 1987 Mar 20;82(3A):11-7.

267 Hajjar IM, Grim CE, Kotchen TA. Dietary calcium lowers the age-related rise in blood pressure in the United States: the NHANES III survey. *J Clin Hypertens (Greenwich)*. 2003 Mar-Apr;5(2):122-6.

268 Temple NJ. Refined carbohydrates - a cause of suboptimal nutrient intake. *Med Hypotheses*. 1983 Apr;10(4):411-24.

269 Aliaibi H. A deleterious interaction between copper deficiency and sugar ingestion may be the missing link in heart disease. *Med Hypotheses*. 2008;70(6):1163-6.

270 Johnson S. The multifaceted and widespread pathology of magnesium deficiency. *Med Hypotheses*. 2001 Feb;56(2):163-70.

271 Sakata K, Matsumura Y, Yoshimura N, Tamaki J, Hashimoto T, Oguri S, Okayama A, Yanagawa H. Relationship between skipping breakfast and cardiovascular disease risk factors in the national nutrition survey data. *Nippon Koshu Eisei Zasshi*. 2001 Oct;48(10):837-41.

272 Caster WO, Parthenides MD, Growthheart, cholesterol, and blood pressure observed in rats fed common breakfast cereals. *Am J Clin Nutr*. 1976 May;29(5):529-34.

273 Ascherio A, Hennekens C, Willett WC, Sacks F, Sato B, Rimm B, Witztman J, Stampfer MJ. Prospective study of nutritional factors, blood pressure, and hypertension among US women. *Hypertension*. 1996 May;27(5):1065-72.

274 Alonso A, Beunza JJ, Bes-Rastrollo M, Pajares RM, Martínez-González MA. Vegetable protein and fiber from cereal are inversely associated with the risk of hypertension in a Spanish cohort. *Arch Med Res*. 2006 Aug;37(6):778-86.

275 Anderson JW. Plant fiber and blood pressure. *Ann Intern Med*. 1983 May;98(5 Pt 2):842-6.

276 Miller WL, Crabtree BF, Evans DK. Exploratory study of the relationship between hypertension and diet diversity among Basa Islanders. *Public Health Rep*. 1992 Jul-Aug;107(4):426-32.

- 285 Djousse L, Rudich T, Gaziano JM. Nut consumption and risk of hypertension in US male physicians. *Clin Nutr*. 2009 Feb;28(1):10-4.
- 286 Welty FK, Lee KS, Lew NS, Zhou JR. Effect of soy nuts on blood pressure and lipid levels in hypertensive, prehypertensive, and normotensive postmenopausal women. *Arch Intern Med*. 2007 May 28;167(10):1060-7.
- 287 Block G, Jensen CD, Norkus EP, Hudes M, Crawford PB. Vitamin C in plasma is inversely related to blood pressure and change in blood pressure during the previous year in young Black and White women. *Nutr J*. 2008 Dec 17;7:35.
- 288 Kim MK, Sasaki S, Sasazuki S, Okubo S, Hayashi M, Tsugane S. Lack of long-term effect of vitamin C supplementation on blood pressure. *Hypertension*. 2002 Dec;40(6):797-803.
- 289 Hsieh YC, Hung CT, Lien LM, Bai CH, Chen WH, Yeh CY, Chen YH, Hsieh FI, Chiu HC, Chiou HY, Hsu CY. A significant decrease in blood pressure through a family-based nutrition health education programme among community residents in Taiwan. *Public Health Nutr*. 2009 Apr;12(4):570-7.
- 290 Shintani TT, Beckham S, Brown AC, O'Connor HK. The Hawaii Diet: a low fat high carbohydrate, low fat multi-cultural diet for the reduction of chronic disease risk factors: obesity, hypertension, hypercholesterolemia, and hyperglycemia. *Hawaii Med J*. 2001 Mar;60(3):69-73.
- 291 Douglass JM, Rasgon IM, Fleiss PM, Schmidt RD, Peters SN, Abelmann EA. Effects of a raw food diet on hypertension and obesity. *South Med J*. 1985 Jul;78(7):841-4.
- 292 Genesis 1:29; 3:18, King James Version of the Holy Bible.
- 293 White. EG. *Counsels on Diet and Foods*, Hagerstown Maryland, Review & Herald Publishing association (1938). http://wordoftruth.seedoftruth.net/downloads/counsels_on_diet_and_foods
- 294 Proverbs 23:2, King James Version of the Holy Bible.
- 295 Antic V, Dulloo A, Montani JP. Short-term (5-day) changes in food intake alter daily hemodynamics in rabbits. *Am J Hypertens*. 2003 Apr;16(4):302-6.
- 296 Ren J. Leptin and hyperleptinemia - from friend to foe for cardiovascular function. *J Endocrinol*. 2004 Apr;181(1):1-10.
- 297 Ecclesiastes 10:17 King James Version of the Holy Bible.
- 298 Dolinsky VW, Morton JS, Oka T, Robillard-Frayne I, Bagdan M, Lopaschuk GD, Des Rosiers C, Walsh K, Davidge ST, Dyck JR. Calorie restriction prevents hypertension and cardiac hypertrophy in the spontaneously hypertensive rat. *Hypertension*. 2010 Sep;56(3):412-21.
- 299 Sharifi AM, Mohseni S, Nekoparvar S, Larjani B, Fakhrzadeh H, Oryan S. Effect of caloric restriction on nitric oxide production, ACE activity, and blood pressure regulation in rats. *Acta Physiol Hung*. 2008 Mar;95(1):55-63.
- 300 McCarty MF. A preliminary fast may potentiate response to a subsequent low-salt, low-fat vegan diet in the management of hypertension - fasting as a strategy for breaking metabolic vicious cycles. *Med Hypotheses*. 2003 May;60(5):624-33.
- 301 Goldhamer AC, Lisle DJ, Sultana P, Anderson SV, Parpia B, Hughes B, Campbell TC. Medically supervised water-only fasting in the treatment of borderline hypertension. *J Altern Complement Med*. 2002 Oct;8(5):643-50.
- 302 Andersson B, Wallin G, Hedner T, Ahlberg AC, Andersson OK. Acute effects of short-term fasting on blood pressure, circulating noradrenaline and efferent sympathetic nerve activity. *Acta Med Scand*. 1988;223(6):485-90.
- 303 Gharbi N, Mornagu B, El-Fazaas S, Kamoun A, Gharib C. Effect of dehydration on nitric oxide, corticotropin and vasopressinergic axis in rat. *C R Biol*. 2004 Jan;327(1):12-20.
- 304 Miyake Y, Kuzuya K, Ueno C, Katayama N, Hayakawa T, Tsuge H, Osawa T. Suppressive Effect of Components in Lemon Juice on Blood Pressure in Spontaneously Hypertensive Rats Food Sci. Technol. Int. Tokyo, 4 (1), 29-32, 1998.
- 305 Inoue T, Iseki K, Iseki C, Kinjo K, Ohya Y, Takishita S. Higher heart rate predicts the risk of developing hypertension in a normotensive screened cohort. *Circ J*. 2007 Nov;71(11):1755-60.
- 306 Pivovska A, Piotrowski W, Broda G, Drygas W, Gluszek J, Zdrojewski T, Kozakiewicz K, Stepaniak U, Bandosz P. The relationship between resting heart rate and atherosclerosis risk factors. *Kardiol Pol*. 2008 Oct;66(10):1069-75.
- 307 Tan Y, Gan Q, Knuemper MM. Central alpha-adrenergic receptors and corticotropin releasing factor mediate hemodynamic responses to acute cold stress. *Brain Res*. 2003 Apr 4;968(1):122-9.
- 308 Tomoda F, Takata M, Kagitani S, Kinuno H, Yasumoto K, Tomita S, Inoue H. Different platelet aggregability during mental stress in two stages of essential hypertension. *Am J Hypertens*. 1999 Nov;12(1 Pt 1):1063-70.
- 309 Markovitz JH, Matthews KA, Kannel WB, Cobb JL, D'Agostino RB. Psychological predictors of hypertension in the Framingham Study. Is there tension in hypertension? *JAMA*. 1993 Nov 24;270(20):2439-43.
- 310 Matthew 6:34. King James Version of the Holy Bible.
- 311 Lal N, Ahuja RC, Madhukar. Life events in hypertensive patients. *J Psychosom Res*. 1982;26(4):441-5.
- 312 Guo ZC. The matched case-control study of the risk factors associated with edema-proteinuria hypertension syndrome (EPHS). *Zhonghua Liu Xing Bing Xue Za Zhi*. 1992 Dec;13(6):351-4.
- 313 Gasperin D, Netuveli G, Dias-da-Costa JS, Pattussi MP. Effect of psychological stress on blood pressure increase: a meta-analysis of cohort studies. *Cad Saude Publica*. 2009 Apr;25(4):715-26.
- 314 Everson SA, Goldberg DE, Kaplan GA, Julkunen J, Salonen JT. Anger expression and incident hypertension. *Psychosom Med*. 1998 Nov-Dec;60(6):730-5.
- 315 Player MS, King DE, Maimous AG 3rd, Geesey ME. Psychosocial factors and progression from prehypertension to hypertension or coronary heart disease. *Ann Fam Med*. 2007 Sep-Oct;5(5):403-11.
- 316 Proverbs16:32, King James Version of the Holy Bible.
- 317 Blanchflower DG, Oswald AJ. Hypertension and happiness across nations. *J Health Econ*. 2008 Mar;27(2):218-33.
- 318 Linden W, Lenz JW, Con AH. Individualized stress management for primary hypertension: a randomized trial. *Arch Intern Med*. 2001 Apr 23;161(8):1071-80.
- 319 Theorell T, Emlund N. On physiological effects of positive and negative life changes—a longitudinal study. *J Psychosom Res*. 1993 Sep;37(6):653-9.
- 320 John 8:3. King James Version of The Holy Bible.
- 321 1John 4:18. King James Version of The Holy Bible.
- 322 Luke 12:22-31. King James Version of The Holy Bible.
- 323 Gerin W, Davidson KW, Christenfeld NJ, Goyal T, Schwartz JE. The role of angry rumination and distraction in blood pressure recovery from emotional arousal. *Schwartz Med*. 2006 Jan-Feb;68(1):64-72.
- 324 Philippians 3:13. King James Version of The Holy Bible.
- 325 Gao X, Nelson ME, Tucker KL. Television viewing is associated with prevalence of metabolic syndrome in Hispanic elders. *Diabetes Care*. 2007 Mar;30(3):694-700.
- 326 Halanach H, Safford MM, Kertesz SG, Pletcher MJ, Kim YI, Person SD, Lewis CE, Kiefe CI. Alcohol consumption in young adults and incident hypertension: 20-year follow-up from the Coronary Artery Risk Development in Young Adults Study. *Am J Epidemiol*. 2010 Mar 1;171(5):532-9.
- 327 Himmelstein DU, Thorne D, Warren E, Woolhandler S. Medical bankruptcy in the United States, 2007: results of a national study. *Am J Med*. 2009 Aug;122(8):741-6.
- 328 Siegrist J. Effort-reward imbalance at work and cardiovascular diseases. *Int J Occup Med Environ Health*. 2010;23(3):279-85.
- 329 Vrijkotte TG, van Doornen LJ, de Geus EJ. Effects of work stress on ambulatory blood pressure, heart rate, and heart rate variability. *Hypertension*. 2000 Apr;35(4):880-6.
- 330 Garcia-Verá MP, Sanz J, Espinosa R, Fortun M, Magan I. Differences in emotional personality traits and stress between sustained hypertension and normotension. *Hypertens Res*. 2010 Mar;33(3):203-8.
- 331 Meizick EJ, Matthews KA, Hall M, Kamarck TW, Strolopp JP, Buysse DJ, Owens JF, Reis SE. Low life purpose and high hostility are related to an attenuated decline in nocturnal blood pressure. *Health Psychol*. 2010 Mar;29(2):196-204.
- 332 Holt-Lunstad J, Uchino BN, Smith TW, Hicks A. On the importance of relationship quality: the impact of ambivalence in friendships on cardiovascular functioning. *Ann Behav Med*. 2007 Jun;33(3):278-90.
- 333 Spitzer SB, Labre MM, Ironson GH, Gellman MD, Schneiderman N. The influence of social situations on ambulatory blood pressure. *Psychosom Med*. 1992 Jan-Feb;54(1):79-86.
- 334 Allen K, Blascovich J, Mendes WB. Cardiovascular reactivity and the presence of pets, friends, and spouses: the truth about cats and dogs. *Psychosom Med*. 2002 Sep-Oct;64(5):727-39.
- 335 Zanini CR, Jardim PC, Salgado CM, Nunes MC, Urrêda FL, Carvalho MV, Pereira DA, Jardim Tde S, Souza WK. Music therapy effects on the quality of life and the blood pressure of hypertensive patients. *Arq Bras Cardiol*. 2009 Nov;93(5):534-40.
- 336 Chafin S, Roy M, Gerin W, Christenfeld N. Music can facilitate blood pressure recovery from stress. *Br J Health Psychol*. 2004 Sep;9(Pt 3):393-403.
- 337 Sobngwi E, Mbanya JC, Unwin NC, Porcher R, Kengne AP, Fezeu L, Minkoulou EM, Tournoux C, Gautier JF, Aspray TJ, Alberti K. Exposure over the life course to an urban environment and its relation with obesity, diabetes, and hypertension in rural and urban Cameroon. *Int J Epidemiol*. 2004 Aug;33(4):769-76.
- 338 Mordukhovich I, Wilker E, Suh H, Wright R, Sparrow D, Vokonas PS, Schwartz J. Black carbon exposure, oxidative stress genes, and blood pressure in a repeated-measures study. *Environ Health Perspect*. 2009 Nov;117(11):1767-72.
- 339 Wiecek A, Kokot F. Does industrial environment influence the prevalence of arterial hypertension, plasma cholesterol and uric acid concentration and activity of the renin-aldosterone system? *Przełg Lek*. 1996;53(4):356-9.
- 340 Chang TY, Su TC, Lin SY, Jain RM, Chan CC. Effects of occupational noise exposure on 24-hour ambulatory vascular properties in male workers. *Environ Health Perspect*. 2007 Nov;115(11):1660-4.
- 341 Haralabidis AS, Dimakopoulou K, Vigna-Taglianti F, Giampaolo M, Borgini A, Dudley ML, Pershagen G, Bluhm G, Houthuijs D, Babich W, Velonakis M, Katsouyanni K, Jarup I, HYENA Consortium. Acute effects of night-time noise exposure on blood pressure in populations living near airports. *Eur Heart J*. 2008 Mar;29(5):658-64.
- 342 Psalms 46:10. King James Version of The Holy Bible.
- 343 Belojević GA, Jakovljević BD, Stojanov VI, Slepević VZ, Panošević KZ. Nighttime road-traffic noise and arterial hypertension in an urban population. *Hypertens Res*. 2008 Apr;31(4):775-81.
- 344 Hartig T, Evans GW, Jamner LD, Davis DS, Garlinge T. Tracking restoration in rural and urban field settings. *J Environ Psych* 2003 23(2):109-23.
- 345 Lohr VI, Pearson-Mims CH, Goodwin GK. Interior Plants May Improve Worker Productivity and Reduce Stress in a Windowless Environment. *J. Environ. Hort*. 14(2):97-100, June 1996
- 346 Park SH, Mattson RH. Ornamental indoor plants in hospital rooms enhanced health outcomes of patients recovering from surgery. *J Altern Complement Med*. 2009 Sep;15(9):975-80.
- 347 Burr JA, Tavares J, Mutcher JE. Volunteering and Hypertension Risk in Later Life. *J Aging Health*. 2011 Feb;23(1):24-51.
- 348 Acts 20:35. King James Version of The Holy Bible.
- 349 Sullivan PA, Procci WR, DeQuattro V, Schoentgen S, Levine D, van der Meulen J, Bornheimer JF. Anger, anxiety, guilt and increased basal and stress-induced neurogenic tone: causes or effects in primary hypertension? *Clin Sci (Lond)*. 1981 Dec;61 Suppl 7:389s-392s.
- 350 Jamner LD, Shapiro D, Hui KK, Oakley ME, Lovett M. Hostility and differences between clinic, self-determined, and ambulatory blood pressure. *Psychosom Med*. 1993 Mar-Apr;55(2):203-11.
- 351 Chafin S, Roy M, Gerin W, Christenfeld N. Music can facilitate blood pressure recovery from stress. *Br J Health Psychol*. 2004 Sep;9(Pt 3):393-403.
- 352 Exodus ch. 20, Deuteronomy ch. 5. King James Version of The Holy Bible.
- 353 Romans 3:20. King James Version of The Holy Bible.
- 354 1Peter 2:22. King James Version of The Holy Bible.
- 355 Romans 3:23. King James Version of The Holy Bible.
- 356 1John 1:8. King James Version of The Holy Bible.
- 357 White EG, Steps to Christ p. 49 <http://www.ted-adventist.org/sites/default/files/Steps%20to%20Christ-EGW.pdf>
- 358 1Timothy 6:12. King James Version of The Holy Bible.
- 359 Philippians 3:14. King James Version of The Holy Bible.
- 360 Romans 2:7. King James Version of The Holy Bible.
- 361 Leviticus 19:18. King James Version of The Holy Bible.
- 362 Matthew 18:15-18. King James Version of The Holy Bible.
- 363 Yan LL, Liu K, Matthews KA, Daviglus ML, Ferguson TF, Kiefe CI. Psychosocial factors and risk of hypertension: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *JAMA*. 2003 Oct 22;290(16):2138-48.
- 364 Revelation 14:12. King James Version of The Holy Bible.
- 365 Lawler-Row KA, Karremans JC, Scott C, Edlis-Mattityahou M, Edwards L. Forgiveness, physiological reactivity and health: the role of anger. *Int J Psychophysiol*. 2008 Apr;68(1):51-8.
- 366 Ephesians 4:32. The King James Version of The Holy Bible.
- 367 Belding JN, Howard MG, McGuire AM, Schwartz AC, Wilson JH. Social buffering by God: prayer and measures of stress. *J Relig Health*. 2010 Jun;49(2):179-87.
- 368 Gillum RF, Ingram DD. Frequency of attendance at religious services, hypertension, and blood pressure: the Third National Health and Nutrition Examination Survey. *Psychosom Med*. 2006 May-Jun;68(3):382-5.
- 369 Koenig HG, George LK, Hays JC, Larson DB, Cohen HJ, Blazer DG. The relationship between religious activities and blood pressure in older adults. *Int J Psychiatry Med*. 1998;28(2):189-213.
- 370 Pickering TG. Mental stress as a causal factor in the development of hypertension and cardiovascular disease. *Curr Hypertens Rep*. 2001 Jun;3(3):249-54.
- 371 Mark 13:8. King James Version of The Holy Bible.
- 372 Krause N, Liang J, Shaw BA, Sugisawa H, Kim HK, Sugihara Y. Religion, death of a loved one, and hypertension among older adults in Japan. *J Gerontol B Psychol Sci Soc Sci*. 2002 Mar;57(2):S96-S107.
- 373 John 14:1-3. The King James Version of The Holy Bible.
- 374 Matthew 16:27. The King James Version of The Holy Bible.
- 375 2Timothy 4:1,8. The King James Version of The Holy Bible.
- 376 Daniel 7. The King James Version of The Holy Bible.
- 377 John 5:28,29. The King James Version of The Holy Bible.
- 378 Psalms 146:4. The King James Version of The Holy Bible.
- 379 Psalms 6:5; 115:17. The King James Version of The Holy Bible.
- 380 Ecclesiastes 9:5; 6. 10. The King James Version of The Holy Bible.
- 381 1Thessalonians 4:15-18. The King James Version of The Holy Bible.
- 382 Job 14:14. The King James Version of The Holy Bible.
- 383 1Corinthians 15:52-58. The King James Version of The Holy Bible.
- 384 Daniel 12:2. The King James Version of The Holy Bible.
- 385 Matt 25:41. The King James Version of The Holy Bible.
- 386 2Peter 3:9. The King James Version of The Holy Bible.
- 387 Malachi 4:1,3. The King James Version of The Holy Bible.
- 388 Matt 3:12. The King James Version of The Holy Bible.
- 389 Jude 1:7. The King James Version of The Holy Bible.
- 390 Revelation 21:4. The King James Version of The Holy Bible.
- 391 Isaiah 65:17. The King James Version of The Holy Bible.
- 392 Nahum 1:9. The King James Version of The Holy Bible.
- 393 Obadiah 1:15,16. The King James Version of The Holy Bible.
- 394 Revelation 22:2. The King James Version of The Holy Bible.